arXiv:1110.2937v1 [math.QA] 13 Oct 2011

PARAMETRIZATIONS OF CANONICAL BASES AND IRREDUCIBLE COMPONENTS OF NILPOTENT VARIETIES

YONG JIANG

Abstract. It is known that the set of irreducible components of nilpotent varieties provides a geometric realization of the crystal basis. For each reduced expression of a Weyl group element, Geiß, Leclerc and Schröer has recently given a parametrization of a subset of irreducible components in studying the cluster structure of the coordinate ring of the corresponding unipotent subgroup. In this paper we show that their parametrization is compatible with Lusztig's parametrization of canonical basis. We also give some interpretations of Lusztig's transition map in terms of irreducible components of nilpotent varieties.

1. Introduction

1.1. Let $\mathfrak{g}$ be a Kac-Moody algebra associated with a symmetric Cartan matrix and $U_q(\mathfrak{g})$ be its quantized enveloping algebra. Denote by $\mathscr{B}(\infty)$ the crystal basis [10] of the negative part $U_q^-(\mathfrak{g})$. It is known that the globalization of crystal basis coincides with Lusztig's canonical basis [14][15].

If $\mathfrak{g}$ is a finite-dimensional simple Lie algebra, the canonical basis of $U_q^-(\mathfrak{g})$ has a labeling of r-tuple of non-negative integers through the PBW-basis [14], where r is the length of the longest element w_0 in the Weyl group. More precisely, for each reduced expression $\mathbf{i}$ of w_0, we have a bijection $\psi_{\mathbf{i}} : \mathbb{N}^r \simeq \mathscr{B}(\infty)$, known as Lusztig's parametrization. It was later generalized to any Kac-Moody algebra in the following sense [17]: For each w of length r and a reduced expression $\mathbf{i}$, we have an injective map $\psi_{\mathbf{i}} : \mathbb{N}^r \hookrightarrow \mathscr{B}(\infty)$. The image does not depend on the choice of $\mathbf{i}$ and thus can be denoted by $\mathscr{B}(w)$. It was shown in [13] that $\mathscr{B}(w)$ provides the dual canonical basis of the quantum unipotent subgroup $U_q^-(w)$ associated to w.

There are many interesting applications of Lusztig's parametrizations of canonical bases. For example, criteria of total positivity [2], combinatorial expressions of tensor product multiplicities [6].

1.2. On the other hand, Kashiwara and Saito [12] has given a geometric construction of the crystal $\mathscr{B}(\infty)$ using Lusztig's nilpotent varieties, which are varieties of certain nilpotent representations of the preprojective algebra Λ associated to the Dynkin diagram of $\mathfrak{g}$. The set $\mathcal{B}$ of irreducible components of nilpotent varieties has a crystal structure isomorphic to $\mathscr{B}(\infty)$.

Recently Geiß, Leclerc and Schröer has shown in [9] that a subset (will be denoted by $\mathcal{B}(w)$ in 3.5) of irreducible components in $\mathcal{B}$ provides the dual semicanonical basis [18] of the coordinate ring $\mathbb{C}[N(w)]$ of the unipotent subgroup associated to w. For each reduced expression $\mathbf{i}$ of w, elements in $\mathcal{B}(w)$ are Zariski closures of certain

---

Supported by the Sonderforschungsbereich 701 "Spectral Structures and Topological Methods in Mathematics" in Bielefeld University.

irreducible constructible subsets $\Lambda_{\mathbf{i}}^{\mathbf{a}}$ ($\mathbf{a} \in \mathbb{N}^r$). And $\Lambda_{\mathbf{i}}^{\mathbf{a}}$ is the set of certain Λ-modules filtered by some particular modules $M_{\mathbf{i},k}$ ($1 \leq k \leq r$) with multiplicities $\mathbf{a}$ (see 3.4 and 3.5 for details). In this way we have an $\mathbb{N}^r$-parametrization of $\mathscr{B}(\infty)$. In other words, we have an injective map $\phi_{\mathbf{i}} : \mathbb{N}^r \hookrightarrow \mathcal{B}$ and the image $\mathcal{B}(w)$ only depends on w.

Thus it is quite natural to ask if the subset $\mathcal{B}(w)$ is the geometric counterpart of $\mathscr{B}(w)$, i. e. the image of $\mathscr{B}(w)$ under the crystal isomorphism $\mathscr{B}(\infty) \simeq \mathcal{B}$ is precisely $\mathcal{B}(w)$. Moreover it would be interesting to know the relationship between Lusztig and GLS's parametrizations.

1.3. In this paper we give an affirmative answer to the above question by showing that for any $w \in W$ and any reduced expression $\mathbf{i}$, Lusztig's parametrization coincides with GLS's, if we identify $\mathscr{B}(\infty)$ with its geometric counterpart $\mathcal{B}$ (Theorem 6.1). This result also means that we have a direct description of the isomorphism $\mathscr{B}(\infty) \simeq \mathcal{B}$ restricting to the subset $\mathscr{B}(w)$ (see Remark 6.4 for details). We then give some interpretations of Lusztig's transition map $R_{\mathbf{i}}^{\mathbf{i}'}$ in terms of irreducible components of nilpotent varieties, where $\mathbf{i}$ and $\mathbf{i}'$ are two reduced expressions differing by a 2 or 3-move.

If $\mathfrak{g}$ is a finite-dimensional simple Lie algebra and $\mathbf{i}$ is a Q-adapted reduced expression of the longest element w_0, the result in our main theorem has recently been proved by Baumann and Kamnitzer [5] (see Remark 6.3 for details). The arguments in [5] in fact work for any Kac-Moody algebra $\mathfrak{g}$ and those element $w \in W$ with a reduced expression $\mathbf{i}$ adapted to some quiver Q. However, even in the case that $\mathfrak{g}$ is finite-dimensional, there exist elements $w \in W$ which do not have any adapted reduced expressions. Thus our theorem is new in all cases.

1.4. A key ingredient in the proof of our main result is the application of reflection functors for preprojective algebras introduced in [5], where it is shown that the reflection functors are geometric analogues of certain crystal operators defined by Saito [20] (see 4.3). We prove that all the modules $M_{\mathbf{i},k}$, whose multiplicities yield GLS's parametrization, can be obtained from simple Λ-modules by applying reflection functors (Proposition 5.6). This result plays an important role in proving Theorem 6.1. On the other hand, it can be viewed as a generalization of a classical result that all indecomposable preprojective and preinjective representations of a quiver can be obtained from simple representations (of a quiver possibly with different orientation) by applying BGP-reflection functors [3]. Furthermore, similar results were proved in [1], where the modules $M_{\mathbf{i},k}$ were introduced differently in a dual form and some derived reflection functors were used.

After a preliminary version of this paper was written, we are informed by Jan Schröer that Benjamin Bolten also introduced reflection functors, very similar to Baumann-Kamnitzer ones, and proved the fact that the modules $M_{\mathbf{i},k}$ can be obtained via reflection functors in his Diploma thesis in June, 2010.

1.5. The paper is organized as follows: In Section 2 we collect basic definitions and properties of quantum groups, crystal basis and Lusztig's parametrization. We then recall the notion of preprojective algebras and nilpotent varieties, also the geometric construction of crystal bases in Section 3. Preliminaries on reflection functors for preprojective algebras are collected in Section 4. The aim of Section 5 is to prove the key result that the modules $M_{\mathbf{i},k}$ can be obtained from simple modules by applying reflection functors (Proposition 5.6). In Section 6 we prove the main results in this paper, namely the compatibility of Lusztig's parametrization of canonical basis and GLS's parametrization of irreducible components of nilpotent varieties (Theorem 6.1). Finally we give some interpretations of Lusztig's transition map $R_{\mathbf{i}}^{\mathbf{i}'}$ in Section 7.

2. Quantum groups and parametrizations of canonical bases

2.1. **Quantum groups.** Let $C = (c_{ij})_{i,j\in I}$ be a symmetric generalized Cartan matrix and $\mathfrak{g}$ be the associated Kac-Moody algebra, where the indexing set $I = \{1, 2, \dots, n\}$. Denote by α_i (resp. h_i) the simple roots (resp. coroots). So $c_{ij} = \langle h_i, \alpha_j\rangle$. Let P be the weight lattice and Q be the root lattice. The i-th fundamental weight is denoted by ϖ_i $(i \in I)$. Let W be the corresponding Weyl group generated by simple reflections s_i $(i \in I)$. The length of an element $w \in W$ is denoted by $\ell(w)$. If $w = s_{i_r} s_{i_{r-1}} \cdots s_{i_1}$ is a reduced expression we say $\mathbf{i} = (i_r, i_{r-1} \dots, i_1)$ is a reduced expression of w.

Let $U_q(\mathfrak{g})$ be the quantized enveloping algebra of $\mathfrak{g}$, which is a $\mathbb{Q}(q)$-algebra with generators e_i, f_i $(i \in I)$ and q^h $(h \in P^*)$. Let $U_q^-(\mathfrak{g})$ be the subalgebra of $U_q(\mathfrak{g})$ generated by f_i $(i \in I)$. The defining relations of $U_q^-(\mathfrak{g})$ are the quantum Serre relations:

$$\sum_{r+s=1-c_{ij}} (-1)^r f_i^{(r)} f_j f_i^{(s)} = 0, \quad (i, j \in I,\ i \neq j).$$

where

$$f_i^{(r)} = f_i^r/[r]!,\ [r]! = [r][r-1]\cdots[1],\ [k] = \frac{q^k - q^{-k}}{q - q^{-1}}.$$

Let $Q_- = -\sum_{i\in I} \mathbb{N}\alpha_i$ be the negative root lattice. The algebra $U_q^-(\mathfrak{g})$ is Q_--graded. That is, for any $\alpha = -\sum_{i\in I} a_i\alpha_i \in Q$, the homogeneous part $U_q^-(\mathfrak{g})_\alpha$ is the $\mathbb{Q}$-space spanned by $f_{i_1} f_{i_2} \cdots f_{i_k}$ where $\sharp\{k|i_k = i\} = a_i$ for all $i \in I$. For a homogeneous element $x \in U_q^-(\mathfrak{g})_\alpha$ we will write $\mathrm{wt}(x) = \alpha$.

2.2. **Quantum unipotent subgroups.** For each $i \in I$, Lusztig defined $\mathbb{Q}(q)$-algebra automorphisms T_i, T_i^{-1} of $U_q(\mathfrak{g})$. We recall the definition of T_i^{-1} here:

$$\begin{aligned}
T_i^{-1}(q^h) &= q^{s_i(h)}, \quad (h \in P^*),\\
T_i^{-1}(e_i) &= -q^{-h_i} f_i,\\
T_i^{-1}(f_i) &= -e_i q^{h_i},\\
T_i^{-1}(e_j) &= \sum_{r+s=-c_{ij}} (-1)^r q^{-r} e_i^{(r)} e_j e_i^{(s)}, \quad (j \neq i),\\
T_i^{-1}(f_j) &= \sum_{r+s=-c_{ij}} (-1)^r q^{r} f_i^{(r)} f_j f_i^{(s)}, \quad (j \neq i),
\end{aligned}$$

Note that T_i (resp. T_i^{-1}) was denoted by $T''_{i,1}$ (resp. $T'_{i,-1}$) in [16].

Now for any $w \in W$ and a reduced expression $\mathbf{i} = (i_r, i_{r-1}, \dots, i_1)$, set

$$\beta_{\mathbf{i},k} = s_{i_1} \cdots s_{i_{k-1}}(\alpha_{i_k}), \quad \text{for any } 1 \leq k \leq r.$$

We know that the set $\Delta_w = \{\beta_{\mathbf{i},k} | 1 \leq k \leq r\}$ consists of all positive roots α such that $w(\alpha)$ is a negative root (Thus it only depends on w).

We then define the following quantum root vectors:

$$F(\beta_{\mathbf{i},k}) = T_{i_1}^{-1} \cdots T_{i_{k-1}}^{-1}(f_{i_k}), \quad \text{for all } 1 \leq k \leq r.$$

Note that these elements are in $U_q^-(\mathfrak{g})$. For $\mathbf{a} = (a_1, \dots, a_r) \in \mathbb{N}^r$ set

$$F_{\mathbf{i}}(\mathbf{a}) = F(\beta_{\mathbf{i},r})^{(a_r)} \cdots F(\beta_{\mathbf{i},1})^{(a_1)}.$$

The subspace of $U_q^-(\mathfrak{g})$ spanned by $\{F_{\mathbf{i}}(\mathbf{a}) | \mathbf{a} \in \mathbb{N}^r\}$ is a subalgebra of $U_q^-(\mathfrak{g})$ and independent of the choice of the reduced expression $\mathbf{i}$ of w. We denote it by $U_q^-(w)$, called the *quantum unipotent subgroup* associated with w (see [13]).

Moreover, the set

$$\mathcal{P}_{\mathbf{i}} = \{F_{\mathbf{i}}(\mathbf{a}) | \mathbf{a} \in \mathbb{N}^r\}$$

is a basis of $U_q^-(w)$, called the PBW-basis attached to $\mathbf{i}$. In particular, if $\mathfrak{g}$ is a finite dimensional simple Lie algebra and $w = w_0$ the longest element in the Weyl group, we have $U_q^-(w_0) = U_q^-(\mathfrak{g})$ and $\mathcal{P}_{\mathbf{i}}$ is a basis of $U_q^-(\mathfrak{g})$.

2.3. **Crystal and canonical bases.** In this subsection we briefly recall the definition of crystal basis following [10].

Let ${}_ir$ be the $\mathbb{Q}(q)$-linear endomorphism of $U_q^-(\mathfrak{g})$ defined by ${}_ir(f_j) = \delta_{ij}$ and

$${}_ir(xy) = {}_ir(x)y + q^{\langle h_i, \alpha \rangle} x {}_ir(y), \quad (x \in U_q^-(\mathfrak{g})_\alpha,\ \alpha \in Q_-,\ y \in U_q^-(\mathfrak{g})).$$

Then there exists a unique non-degenerate symmetric bilinear form such that $(1, 1) = 1$ and

$$(x, f_i y) = ({}_ir(x), y), \quad \text{for} \quad x, y \in U_q^-(\mathfrak{g}), i \in I.$$

Each element $x \in U_q^-(\mathfrak{g})$ can be uniquely written as

$$x = \sum_{k \geq 0} f_i^{(k)} x_k,$$

where ${}_ir(x_k) = 0$ for any $k \geq 0$.

Then the Kashiwara operators $\tilde{e}_i$ and $\tilde{f}_i$ are defined by

$$\tilde{e}_i x = \sum_{k \geq 1} f_i^{(k-1)} x_k, \quad \tilde{f}_i x = \sum_{k \geq 0} f_i^{(k+1)} x_k.$$

Let $\mathcal{A}_0$ be the subring of $\mathbb{Q}(q)$ consisting of all functions having no pole at $q = 0$. Let $\mathscr{L}(\infty)$ be the $\mathcal{A}_0$-submodule of $U_q^-(\mathfrak{g})$ generated by $\tilde{f}_{i_1} \cdots \tilde{f}_{i_l} 1$ ($l \geq 0$, $i_1, \dots, i_l \in I$). Let $\mathscr{B}(\infty)$ be the subset of $\mathscr{L}(\infty)/q\mathscr{L}(\infty)$ consisting of $\tilde{f}_{i_1} \cdots \tilde{f}_{i_l} 1$ (mod $q\mathscr{L}(\infty)$). Then the pair $(\mathscr{L}(\infty), \mathscr{B}(\infty))$ has the following properties:

(1). $\mathscr{L}(\infty)$ is a free $\mathcal{A}_0$-module such that $\mathbb{Q}(q) \otimes_{\mathcal{A}_0} \mathscr{L}(\infty) \cong U_q^-(\mathfrak{g})$.

(2). $\mathscr{B}(\infty)$ is a basis of the $\mathbb{Q}$-vector space $\mathscr{L}(\infty)/q\mathscr{L}(\infty)$.

(3). $\tilde{e}_i \mathscr{L}(\infty) \subset \mathscr{L}(\infty)$ and $\tilde{f}_i \mathscr{L}(\infty) \subset \mathscr{L}(\infty)$ for any $i \in I$ (Hence $\tilde{e}_i$ and $\tilde{f}_i$ act on $\mathscr{L}(\infty)/q\mathscr{L}(\infty)$).

(4). $\tilde{e}_i \mathscr{B}(\infty) \subset \mathscr{B}(\infty) \cup \{0\}$ and $\tilde{f}_i \mathscr{B}(\infty) \subset \mathscr{B}(\infty)$.

(5). For any $b \in \mathscr{B}(\infty)$, $\tilde{e}_i b \in \mathscr{B}(\infty)$ implies $b = \tilde{f}_i \tilde{e}_i b$.

The pair $(\mathscr{L}(\infty), \mathscr{B}(\infty))$ (or just $\mathscr{B}(\infty)$, for simplicity) is called the *crystal basis* of $U_q^-(\mathfrak{g})$. It can be globalized to a basis of $U_q^-(\mathfrak{g})$, which coincides with the *canonical basis* introduced by Lusztig [14][15], denoted by $\mathbf{B}$. The globalization map is denoted by $G : \mathscr{B}(\infty) \to \mathbf{B}$, assigning each $b \in \mathscr{B}(\infty)$ to the corresponding canonical basis $G(b) \in \mathbf{B}$.

2.4. **Abstract crystals.** In [11] Kashiwara introduced the notion of (abstract) crystals. A *crystal* associated with $\mathfrak{g}$ is a set $\mathscr{B}$ endowed with maps $\mathrm{wt} : \mathscr{B} \to P$, $\varepsilon_i, \varphi_i : \mathscr{B} \to \mathbb{Z} \cup \{-\infty\}$ and $\tilde{e}_i, \tilde{f}_i : \mathscr{B} \to \mathscr{B} \cup \{0\}$ (for each $i \in I$) satisfying the following properties:

(1). $\varphi_i(b) = \varepsilon_i(b) + \langle h_i, \mathrm{wt}(b)\rangle$.

(2). If $\tilde{e}_i b \in \mathscr{B}$, then $\mathrm{wt}(\tilde{e}_i b) = \mathrm{wt}(b) + \alpha_i$, $\varepsilon_i(\tilde{e}_i b) = \varepsilon_i(b) - 1$ and $\varphi_i(\tilde{e}_i b) = \varphi_i(b) + 1$.

(3). If $\tilde{f}_i b \in \mathscr{B}$, then $\mathrm{wt}(\tilde{f}_i b) = \mathrm{wt}(b) - \alpha_i$, $\varepsilon_i(\tilde{f}_i b) = \varepsilon_i(b) + 1$ and $\varphi_i(\tilde{f}_i b) = \varphi_i(b) - 1$.

(4). $b' = \tilde{e}_i b$ if and only if $b = \tilde{f}_i b'$ for all $b, b' \in \mathscr{B}$ and $i \in I$.

(5). If $\varphi_i(b) = -\infty$, then $\tilde{e}_i b = \tilde{f}_i b = 0$.

The crystal basis $\mathscr{B}(\infty)$ is an example of a crystal: The maps $\mathrm{wt}, \widetilde{e}_i, \widetilde{f}_i$ have already been defined. For any $b \in \mathscr{B}(\infty)$, we set $\varepsilon_i(b) = \max\{k \in \mathbb{N} | \tilde{e}_i^k(b) \neq 0\}$ and $\varphi_i(b) = \varepsilon_i(b) + \langle h_i, \mathrm{wt}(b)\rangle$.

For two crystals $\mathscr{B}_1$ and $\mathscr{B}_2$, a *morphism* Ψ from $\mathscr{B}_1$ to $\mathscr{B}_2$ is a map $\mathscr{B}_1 \cup \{0\} \mapsto \mathscr{B}_2 \cup \{0\}$ satisfying the following conditions:

(1). $\Psi(0) = 0$.

(2). If $b \in \mathscr{B}_1$ and $\Psi(b) \in \mathscr{B}_2$, then $\mathrm{wt}(\Psi(b)) = \mathrm{wt}(b)$, $\varepsilon_i(\Psi(b)) = \varepsilon_i(b)$ and $\varphi(\Psi(b)) = \varphi(b)$.

(3). If $b, b' \in \mathscr{B}_1$, $\Psi(b), \Psi(b') \in \mathscr{B}_2$ and $\widetilde{f}_i(b) = b'$, then $\widetilde{f}_i(\Psi(b)) = \Psi(b')$.

A morphism Ψ is called an *isomorphism* if it is a bijection and commutes with all $\widetilde{e}_i$ and $\widetilde{f}_i$.

2.5. **The $*$-crystal structure.** We have a $\mathbb{Q}(q)$-algebra anti-involution $* : U_q(\mathfrak{g}) \to U_q(\mathfrak{g})$ defined by

$$*(e_i) = e_i, \quad *(f_i) = f_i, \quad *(q^h) = q^{-h}.$$

It has been proved in [10][11] that

$$*(\mathscr{L}(\infty)) = \mathscr{L}(\infty), \quad *(\mathscr{B}(\infty)) = \mathscr{B}(\infty).$$

Now for each $i \in I$ we define two operators $\widetilde{e}_i^*$ and $\widetilde{f}_i^*$ on $\mathscr{B}(\infty)$ as follows:

$$\widetilde{e}_i^*(b) = (* \circ \widetilde{e}_i \circ *)(b), \quad \widetilde{f}_i^*(b) = (* \circ \widetilde{f}_i \circ *)(b).$$

We also define two maps $\varepsilon_i^*, \varphi_i^* : \mathscr{B}(\infty) \to \mathbb{Z}$ by

$$\varepsilon_i^*(b) = \max\{k \in \mathbb{N} | \widetilde{e}_i^{*k}(b) \neq 0\}, \quad \varphi_i^*(b) = \varepsilon_i^*(b) + \langle h_i, \mathrm{wt}(b)\rangle.$$

Then $(\mathscr{B}(\infty); \mathrm{wt}, \widetilde{e}_i^*, \widetilde{f}_i^*, \varepsilon_i^*, \varphi_i^*)$ is also a crystal. Note that this crystal structure is different from $(\mathscr{B}(\infty); \mathrm{wt}, \widetilde{e}_i, \widetilde{f}_i, \varepsilon_i, \varphi_i)$.

We will use the following notations: For any $b \in \mathscr{B}(\infty)$, write $\widetilde{e}_i^{\max} b = \widetilde{e}_i^{\varepsilon_i(b)} b$ and $\widetilde{e}_i^{*\max} b = \widetilde{e}_i^{*\varepsilon_i^*(b)} b$.

2.6. **The operators $\mathcal{T}_i$.** In [20], Saito introduced operators $\mathcal{T}_i$ and $\mathcal{T}_i^{-1}$ (originally denoted by Λ_i and Λ_i^{-1}) on $\mathscr{B}(\infty)$ as follows

$$\mathcal{T}_i(b) = \widetilde{f}_i^{*\varphi_i(b)} \widetilde{e}_i^{\max} b, \quad \mathcal{T}_i^{-1}(b) = \widetilde{f}_i^{\varphi_i^*(b)} \widetilde{e}_i^{*\max} b.$$

These operators set up a bijection between the following two subsets of $\mathscr{B}(\infty)$:

$$\begin{tikzcd} \{b \in \mathscr{B}(\infty) | \varepsilon_i^*(b) = 0\} \arrow[r, shift left, "\mathcal{T}_i"] & \{b \in \mathscr{B}(\infty) | \varepsilon_i(b) = 0\} \arrow[l, shift left, "\mathcal{T}_i^{-1}"] \end{tikzcd}$$

Moreover, they are analogues of Lusztig's automorphism T_i, T_i^{-1} on the level of crystals.

Proposition 2.1 ([20])**.** *For any* $b \in \mathscr{B}(\infty)$ *with* $\varepsilon_i^*(b) = 0$*, we have*

$$T_i(G(b)) \equiv \mathcal{T}_i b \mod q\mathscr{L}(\infty).$$

Note that in general T_i does not restrict to an automorphism of $U_q^-(\mathfrak{g})$. But for $b \in \mathscr{B}(\infty)$ with $\varepsilon_i^*(b) = 0$, $T_i(G(b))$ lies in $U_q^-(\mathfrak{g})$.

2.7. The Lusztig parametrization of $\mathscr{B}(w)$.

We have the following result:

Theorem 2.2 ([17])**.** *Let* $w \in W$ *and* $\mathbf{i}$ *be a reduced expression of* w*.*

(1). For any $\mathbf{a} = (a_1, \dots, a_r) \in \mathbb{N}^r$*, there exists a unique element* $b_{\mathbf{i},\mathbf{a}} \in \mathscr{B}(\infty)$ *such that*

$$F_{\mathbf{i}}(\mathbf{a}) \equiv b_{\mathbf{i},\mathbf{a}} \mod q\mathscr{L}(\infty).$$

(2). The map $\psi_{\mathbf{i}} : \mathbb{N}^r \to \mathscr{B}(\infty)$ *defined by* $\mathbf{a} \mapsto b_{\mathbf{i},\mathbf{a}}$ *is injective and the image does not depend on the choice of* $\mathbf{i}$*.*

Thus we can denote by $\mathscr{B}(w)$ the image of the map. The theorem gives a $\mathbb{N}^r$-parametrization of elements in $\mathscr{B}(w)$. For each $b \in \mathscr{B}(w)$, $\psi_{\mathbf{i}}^{-1}(b)$ is called the $\mathbf{i}$-*Lusztig data* of b.

Note that when $\mathfrak{g}$ is a finite-dimensional simple Lie algebra and $w = w_0$ the longest element in the Weyl group, the map $\mathbf{a} \mapsto b_{\mathbf{i},\mathbf{a}}$ is a bijection and we have $\mathscr{B}(w) = \mathscr{B}(\infty)$.

3. PREPROJECTIVE ALGEBRAS AND NILPOTENT VARIETIES

3.1. Preprojective algebras.

Let $Q = (I, \Omega)$ be a finite connected quiver without oriented cycles, where I is the vertex set and Ω the arrow set. As in section 2, we set $I = \{1, 2, \dots, n\}$. Denote by s and t the source and target maps from Ω to I. We can associate with Q a symmetric Cartan matrix $C = (c_{ij})_{i,j\in I}$ as follows:

$$c_{ij} = \begin{cases} 2 & \text{if } i = j \\ -\sharp\{a \in \Omega | s(a) = i, t(a) = j; \text{or } s(a) = j, t(a) = i\} & \text{if } i \neq j \end{cases}$$

Thus we can associate a Kac-Moody algebra $\mathfrak{g} = \mathfrak{g}(Q)$ to each quiver Q. The underlying graph of Q is just the Dynkin graph of $\mathfrak{g}$.

Let $\mathbb{C}Q$ be the path algebra of Q. A $\mathbb{C}Q$-module (or a $\mathbb{C}$-representation of Q) M consists of an I-graded $\mathbb{C}$-vector space $M = \oplus_{i\in I} M_i$ and a collection of linear maps $M_a : M_{s(a)} \to M_{t(a)}$ for each arrow $a \in \Omega$.

The *double quiver* $\overline{Q}$ is obtained from Q by adding to each arrow $a : i \to j$ an opposite arrow $a^* : j \to i$. Thus the arrow set of $\overline{Q}$ is $H = \Omega \sqcup \Omega^*$. We extend $*$ to H by setting $(a^*)^* = a$ for each $a \in \Omega$. We also define a map $\epsilon : H \to \{\pm 1\}$ by assigning $\epsilon(a) = 1$ if $a \in \Omega$ and $\epsilon(a) = -1$ if $a \in \Omega^*$. Let c be the two-sided ideal of the path algebra $\mathbb{C}\overline{Q}$ generated by $\sum_{a\in H} \epsilon(a)aa^*$. The quotient algebra $\mathbb{C}\overline{Q}/(c)$ is called the *preprojective algebra* of Q, denoted by $\Lambda = \Lambda(Q)$. It is clear that the path algebra $\mathbb{C}Q$ is a subalgebra of Λ.

A Λ-module is a $\mathbb{C}\overline{Q}$-module M satisfying the following relations:

$$\sum_{a\in H, t(a)=i} \epsilon(a) M_a M_{a^*} = 0, \text{ for each } i \in I.$$

Denote by $\mathrm{mod}(\Lambda)$ the category of finite dimensional Λ-modules. For a module $M \in \mathrm{mod}\,(\Lambda)$, the vector $\underline{\dim}\, M = (\dim M_i)_{i\in I} \in \mathbb{N}^n$ is called the *dimension vector* of M. It will be convenient for us to view the dimension vector as an element in the root lattice by setting $\underline{\dim}\, M = \sum_{i\in I}(\dim M_i)\alpha_i \in Q_+$.

For each vertex $i \in I$ we have a one-dimensional simple Λ-module S_i concentrated on the vertex i. In general there exist other finite-dimensional simple modules. A Λ-module M is called *nilpotent* if there is a composition series of M contains only the simple modules S_i $(i \in I)$. Let $\mathrm{nil}(\Lambda)$ be the full subcategory of $\mathrm{mod}\,(\Lambda)$ consisting of finite-dimensional nilpotent Λ-modules. It is an abelian subcategory.

For a Λ-module M, the *i-socle* (resp. *i-top*) of M is the S_i-isotypic component of the socle (resp. top) of M, denoted by $\mathrm{soc}_i\, M$ (resp. $\mathrm{top}_i\, M$).

We will need the following result, which can be seen from [7] Lemma 1.

Lemma 3.1. *For any Λ-modules X, Y, we have*

$$\dim \mathrm{Ext}^1_\Lambda(X,Y) = \dim \mathrm{Ext}^1_\Lambda(Y,X)$$

3.2. **Nilpotent varieties.** Let $\nu = \sum_{i\in I}\nu_i\alpha_i \in Q_+$ be a dimension vector. Define $\mathrm{mod}(\Lambda,\nu)$ to be the affine variety of all linear maps

$$(M_a)_{a\in H} \in \prod_{a\in H} \mathrm{Hom}_{\mathbb{C}}(\mathbb{C}^{\nu_{s(a)}}, \mathbb{C}^{\nu_{t(a)}})$$

satisfying the following relations

$$\sum_{a\in H, t(a)=i} \epsilon(a)M_aM_{a^*} = 0, \text{ for each } i \in I.$$

Thus a point in $\mathrm{mod}(\Lambda,\nu)$ is just a Λ-module M with $M_i = \mathbb{C}^{\nu_i}$ for all $i \in I$.

The group $\mathrm{GL}(\nu) = \prod_{i\in I}\mathrm{GL}_{\nu_i}(\mathbb{C})$ acts on $\mathrm{mod}(\Lambda,\nu)$ by conjugation. The orbits can be identified with isomorphism classes of Λ-modules with dimension vector ν.

Let $\Lambda(\nu)$ be the subset of $\mathrm{mod}\,(\Lambda,\nu)$ consisting of nilpotent Λ-modules. It is not difficult to see that $\Lambda(\nu)$ is also an affine variety. These varieties are usually called *nilpotent varieties.* The $\mathrm{GL}(\nu)$-action on $\mathrm{mod}(\Lambda,\nu)$ restricts to a $\mathrm{GL}(\nu)$-action on $\Lambda(\nu)$.

3.3. **Geometric crystals.** Denote by $\mathrm{Irr}\,\Lambda(\nu)$ the set of irreducible components of the variety $\Lambda(\nu)$. Let $\mathcal{B} = \sqcup_{\nu\in Q_+}\mathrm{Irr}\,\Lambda(\nu)$. Kashiwara and Saito [12] associated a crystal structure on $\mathcal{B}$ and proved that it is isomorphic to $\mathscr{B}(\infty)$ as abstract crystals. We briefly recall their construction here.

For $i \in I$ and $c \in \mathbb{N}$, define

$$\Lambda(\nu)_{i,c} = \{M \in \Lambda(\nu) | \dim \mathrm{top}_i\, M = c\}.$$

They are locally closed subsets and $\Lambda(\nu) = \sqcup_{c\in\mathbb{N}}\Lambda(\nu)_{i,c}$ for each i. Thus for any irreducible component Z of $\Lambda(\nu)$, there is only one value c such that $Z \cap \Lambda(\nu)_{i,c}$ is open dense in Z. We then define two maps $\varepsilon_i(Z) = c$ and $\varphi_i(Z) = \varepsilon_i(Z) + \langle h_i, \mathrm{wt}(Z)\rangle$, where we set $\mathrm{wt}(Z) = -\nu$ for $Z \in \Lambda(\nu)$.

It is also known that the map $Z \mapsto Z \cap \Lambda(\nu)_{i,c}$ gives a bijection

$$\{Z \in \mathrm{Irr}\,\Lambda(\nu) | \varepsilon_i(Z) = c\} \longleftrightarrow \mathrm{Irr}\,\Lambda(\nu)_{i,c}.$$

Denote by $\Omega(\nu, i, c)$ the set of triples (M, N, f) where $M \in \Lambda(\nu)_{i,0}$, $N \in \Lambda(\nu + c\alpha_i)_{i,c}$ and $f : M \to N$ is an injective morphism of Λ-modules. We then have the following diagram

$$\Lambda(\nu)_{i,0} \xleftarrow{p} \Omega(\nu, i, c) \xrightarrow{q} \Lambda(\nu + c\alpha_i)_{i,c},$$

where p and q are the first and second projections. Then p is a locally trivial fibration with a smooth and connected fibre and q is a principal $\mathrm{GL}(\nu)$-bundle. Hence the above diagram induces the following bijection

$$\begin{tikzcd} \{Z \in \operatorname{Irr} \Lambda(\nu) | \varepsilon_i(Z) = 0\} \arrow[r, shift left, "\widetilde{f}_i^c"] & \{Z \in \operatorname{Irr} \Lambda(\nu + c\alpha_i) | \varepsilon_i(Z) = c\} \arrow[l, shift left, "\widetilde{e}_i^{\max}"] \end{tikzcd}$$

Then we define operators $\widetilde{e}_i, \widetilde{f}_i : \mathcal{B} \to \mathcal{B} \cup \{0\}$ as follows:

$$\widetilde{e}_i(Z) = \widetilde{f}_i^{c-1} \widetilde{e}_i^{\max}(Z), \ \widetilde{f}_i(Z) = \widetilde{f}_i^{c+1} \widetilde{e}_i^{\max}(Z), \ \text{for any } Z \text{ with } \varepsilon_i(Z) = c.$$

Theorem 3.2 ([12])**.** *Equipped with the maps* $\mathrm{wt}, \varepsilon_i, \varphi_i, \widetilde{e}_i, \widetilde{f}_i$ *defined above, the set* $\mathcal{B} = \sqcup_{\nu \in Q_+} \operatorname{Irr} \Lambda(\nu)$ *is a crystal. Moreover, there is a (unique) crystal isomorphism* $\Psi : \mathscr{B}(\infty) \to \mathcal{B}$.

We also have a geometric version of the $*$-crystal structure of $\mathscr{B}(\infty)$ defined in 2.5. For any $M \in \mathrm{mod}(\Lambda)$, we have the dual $M^* \in \mathrm{mod}(\Lambda)$ defined by taking dual vector space $(M^*)_i = M_i^*$ and transpose of linear maps $(M^*)_a = M_{a^*}^{\mathrm{tr}}$ for each $i \in I$ and $a \in H$. This gives an involution on the objects of $\mathrm{nil}(\Lambda)$ and keeps the dimension vectors. And it induces an involution on $\operatorname{Irr} \Lambda(\nu)$ for each $\nu \in Q_+$. Therefore we obtain a weight-preserving involution on $\mathcal{B}$, denoted again by $*$.

Then we define $\varepsilon_i^*(Z) = \varepsilon_i(Z^*)$, $\varphi_i^*(Z) = \varphi_i(Z^*)$, $\widetilde{e}_i^* = * \circ \widetilde{e}_i \circ *$, $\widetilde{f}_i^* = * \circ \widetilde{f}_i \circ *$. The set $\mathcal{B}$ together with these maps is a crystal isomorphic to $(\mathscr{B}(\infty); \mathrm{wt}, \varepsilon_i^*, \varphi_i^*, \widetilde{e}_i^*, \widetilde{f}_i^*)$.

3.4. **The subcategory $\mathcal{C}_w$.** In this and the next subsection we recall some results in [9] concerning parametrizations of a subset of irreducible components in $\mathcal{B}$.

For each $i \in I$, let $\widehat{I}_i$ be the indecomposable injective Λ-module with socle S_i. Note that these modules are infinite-dimensional if Q is not a Dynkin quiver.

For a Λ-module X and a simple module S_j ($j \in I$), recall that $\mathrm{soc}_j X$ is the sum of all submodules U of X with $U \cong S_j$ (see 3.1). For a sequence $(j_1, \ldots, j_t)$ of indices with $1 \leq j_p \leq n$ for all p, there is a unique chain

$$0 = X_0 \subseteq X_1 \subseteq \cdots \subseteq X_t \subseteq X$$

of submodules of X such that $X_p / X_{p-1} = \mathrm{soc}_{j_p}(X / X_{p-1})$. Define $\mathrm{soc}_{(j_1, \ldots, j_t)}(X)$ to be the submodule X_t in the above chain.

Let $\mathbf{i} = (i_r, \ldots, i_1)$ be a reduced expression of an element w in the Weyl group. For $1 \leq k \leq r$, define

$$V_{\mathbf{i},k} = \mathrm{soc}_{(i_k, \ldots, i_1)}(\widehat{I}_k)$$

and set $V_{\mathbf{i}} = \oplus_{k=1}^r V_{\mathbf{i},k}$. Let $\mathcal{C}_{\mathbf{i}} = \mathrm{Fac}(V_{\mathbf{i}})$, i.e. the subcategory of all Λ-modules X such that there exists an epimorphism $V_{\mathbf{i}}^m \twoheadrightarrow X$ for some $m \geq 1$. In fact it does not depend on the choice of $\mathbf{i}$. So we can denote it by $\mathcal{C}_w$ (This subcategory was alternatively defined in [4]). It is clear that $\mathcal{C}_w$ is a subcategory of $\mathrm{nil}(\Lambda)$. Note that when Q is a Dynkin quiver and $w = w_0$ is the longest element in the Weyl group, then we have $\mathcal{C}_{w_0} = \mathrm{nil}(\Lambda) = \mathrm{mod}(\Lambda)$.

3.5. **Parametrizations of irreducible components.** We keep the notations in the last subsection. For $1 \leq k \leq r$ we define

$$k^- = \max\{0, 1 \leq s \leq k-1 | i_s = i_k\}.$$

For each $1 \leq k \leq r$ there is a canonical embedding

$$\iota_k : V_{\mathbf{i},k^-} \to V_{\mathbf{i},k},$$

where for $k = 1$ we set $V_{\mathbf{i},0} = 0$. Let $M_{\mathbf{i},k}$ be the cokernel of ι_k and define

$$M_{\mathbf{i}} = \oplus_{k=1}^r M_{\mathbf{i},k}.$$

For $\mathbf{a} = (a_r, \dots, a_1) \in \mathbb{N}^r$, let $\mathcal{C}_{\mathbf{i},\mathbf{a}}$ be the category of all Λ-modules X such that there exists a chain

$$0 = X_0 \subseteq X_1 \subseteq \cdots \subseteq X_r = X$$

of submodules of X such that $X_k/X_{k-1} \cong M_{\mathbf{i},k}^{a_k}$ for all $1 \leq k \leq r$.

It was shown in [9] that the categories $\mathcal{C}_{\mathbf{i},\mathbf{a}}$ ($\mathbf{a} \in \mathbb{N}$) are pairwise disjoint and

$$\bigsqcup_{\mathbf{a} \in \mathbb{N}} \mathcal{C}_{\mathbf{i},\mathbf{a}} = \mathcal{C}_w.$$

By definition all the modules $X \in \mathcal{C}_{\mathbf{i},\mathbf{a}}$ have the same dimension vector $\mu(\mathbf{a}) = \sum_{k=1}^r a_k \, \underline{\dim}_\Lambda(M_k)$. Let Let $\Lambda(\nu)^w = \{X \in \Lambda(\nu) | X \in \mathcal{C}_w\}$ for any $\nu \in Q_+$ and $\Lambda_{\mathbf{i}}^{\mathbf{a}} = \{X \in \Lambda(\mu(\mathbf{a})) | X \in \mathcal{C}_{\mathbf{i},\mathbf{a}}\}$. Then we have a finite decomposition

$$\Lambda(\nu)^w = \bigsqcup_{\mathbf{a} \in \mathbb{N}^r, \mu(\mathbf{a}) = \nu} \Lambda_{\mathbf{i}}^{\mathbf{a}}$$

into disjoint subsets.

Proposition 3.3 ([9])**.** *$\Lambda_{\mathbf{i}}^{\mathbf{a}}$ is an irreducible constructible subset of $\Lambda(\mu(\mathbf{a}))$ and the Zariski closure $Z_{\mathbf{i}}^{\mathbf{a}}$ of $\Lambda_{\mathbf{i}}^{\mathbf{a}}$ is an irreducible component. In particular, $Z_{\mathbf{i}}^{\mathbf{a}}$ is the unique irreducible component of $\Lambda(\mu(\mathbf{a}))$ which contains a dense open subset belonging to $\Lambda_{\mathbf{i}}^{\mathbf{a}}$.*

Thus we have an injective map $\phi_{\mathbf{i}} : \mathbb{N}^r \to \mathcal{B}$ given by $\mathbf{a} \mapsto Z_{\mathbf{i}}^{\mathbf{a}}$. And from [9] we can see that the image of the map $\phi_{\mathbf{i}}$ is independent of the choice of $\mathbf{i}$, since it provides the dual semicanonical basis of the coordinate ring of the unipotent subgroup $N(w)$, which certainly only depends on w. So we can denote the image by $\mathcal{B}(w)$.

4. Reflection functors for preprojective algebras

4.1. **Definition of reflection functors.** We recall the definition of reflection functors for preprojective algebras following [5]. They are natural generalization of BGP-reflection functors for representations of quivers [3].

Let $M \in \operatorname{mod}(\Lambda)$ and fix $i \in I$. It is convenient to break the data of a Λ-module M into two parts. The first part consists of the vector spaces M_j for $j \neq i$ and the linear maps between them. The second part consists of the vector spaces and linear maps appearing in the following diagram

$$\bigoplus_{a \in H, t(a)=i} M_{s(a)} \xrightarrow{(\varepsilon(a) M_a)} M_i \xrightarrow{(M_{a^*})} \bigoplus_{a \in H, t(a)=i} M_{s(a)}.$$

We simply write the above diagram as

$$(*) \qquad \widetilde{M_i} \xrightarrow{M_{\mathrm{in}(i)}} M_i \xrightarrow{M_{\mathrm{out}(i)}} \widetilde{M_i}.$$

Note that we have $M_{\mathrm{in}(i)}M_{\mathrm{out}(i)} = 0$ since M is a Λ-module.

Now we define a Λ-module $\Sigma_i M$ by replacing the diagram $(*)$ with

$$\begin{tikzcd}
\widetilde{M_i} \arrow[r, "\overline{M}_{\mathrm{out}(i)}M_{\mathrm{in}(i)}"] & \ker M_{\mathrm{in}(i)} \arrow[r, hook, "\iota"] & \widetilde{M_i},
\end{tikzcd}$$

where ι is the natural embedding and the map $\overline{M}_{\mathrm{out}(i)}$ is the unique map such that $\iota\overline{M}_{\mathrm{out}(i)} = M_{\mathrm{out}(i)}$. The remaining data of M, i.e. the vector space M_j for $j \neq i$ and the linear maps between them remain unchanged. It is clear that $\iota\overline{M}_{\mathrm{out}(i)}M_{\mathrm{in}(i)} = 0$, thus $\Sigma_i M$ is a Λ-module.

We can also replace $*$ by

$$\begin{tikzcd}
\widetilde{M_i} \arrow[r, two heads, "\rho"] & \operatorname{coker} M_{\mathrm{out}(i)} \arrow[r, "M_{\mathrm{out}(i)}\overline{M}_{\mathrm{in}(i)}"] & \widetilde{M_i},
\end{tikzcd}$$

where the map $\overline{M}_{\mathrm{in}(i)}$ is the unique map such that $\overline{M}_{\mathrm{in}(i)}\rho = M_{\mathrm{in}(i)}$. In this way we also define a Λ-module, denoted by $\Sigma_i^* M$.

The actions of Σ_i and Σ_i^* on morphisms can be defined in a natural way. So we have defined two functors $\Sigma_i, \Sigma_i^* : \mathrm{mod}(\Lambda) \to \mathrm{mod}(\Lambda)$. It is also easy to see that both functors can be restricted on the subcategory $\mathrm{nil}(\Lambda)$.

Remark 4.1. If M is a $\mathbb{C}Q$-module and i is a sink (resp. source) of the quiver Q, we can see that $\Sigma_i M = \Phi_i^+ M$ (resp. $\Sigma_i^* M = \Phi_i^- M$) where $\Phi_i^\pm$ are the BGP-reflection functors for representations of the quiver Q.

4.2. **Properties of reflection functors.**

We collect some basic properties of reflection functors in the following lemma.

Lemma 4.2 ([5])**.** *(1). Σ_i is left exact and Σ_i^* is right exact.*

(2). We have the following functorial short exact sequences:

$$0 \to \mathrm{soc}_i \to \mathrm{id} \to \Sigma_i\Sigma_i^* \to 0,$$

$$0 \to \Sigma_i^*\Sigma_i \to \mathrm{id} \to \mathrm{top}_i \to 0.$$

(3). Let i and j be two vertices of Q such that they are linked by one single arrow. Then the functors $\Sigma_i\Sigma_j\Sigma_i$ and $\Sigma_j\Sigma_i\Sigma_j$ are isomorphic.

Let $\mathrm{mod}(\Lambda)[i]$ (resp. $\mathrm{mod}(\Lambda)[i]^*$) be the subcategory of $\mathrm{mod}(\Lambda)$ consisting of modules with trivial i-top (resp. i-socle). By (2) in the above lemma we can see that Σ_i and Σ_i^* give inverse equivalences of categories

$$\begin{tikzcd}
\mathrm{mod}(\Lambda)[i] \arrow[r, shift left, "\Sigma_i"] & \mathrm{mod}(\Lambda)[i]^* \arrow[l, shift left, "\Sigma_i^*"]
\end{tikzcd}$$

In general the functor Σ_i is not right exact and Σ_i^* is not left exact. But we have the following easy lemma:

Lemma 4.3. *The restriction of Σ_i (resp. Σ_i^*) on $\mathrm{mod}(\Lambda)[i]$ (resp. $\mathrm{mod}(\Lambda)[i]^*$) is an exact functor.*

Proof. We only prove the statement for Σ_i. The one for Σ_i^* can be proved similarly.

It is sufficient to show that if $f : M \to N$ is surjective and M has trivial i-top (so N also has trivial i-top), then the map $\Sigma_i f : \Sigma_i M \to \Sigma_i N$ is also surjective.

For any $j \neq i$, the map $(\Sigma_i f)_j = f_j$ is surjective by assumption. It remains to check $(\Sigma_i f)_i$, which is given by the following commutative diagram:

$$\begin{tikzcd}
\ker(M_{\mathrm{in}(i)}) \arrow[r, "\iota"] \arrow[d, "(\Sigma_i f)_i"] & \widetilde{M}_i \arrow[r, "M_{\mathrm{in}(i)}"] \arrow[d, "(f_{s(a)})_{t(a)=i}"] & M_i \arrow[d, "f_i"] \\
\ker(N_{\mathrm{in}(i)}) \arrow[r, "\iota"] & \widetilde{N}_i \arrow[r, "N_{\mathrm{in}(i)}"] & N_i
\end{tikzcd}$$

Since M and N have trivial i-top, the map $M_{\mathrm{in}(i)}$ and $N_{\mathrm{in}(i)}$ are both surjective, which means the two rows in the above diagram are in fact exact sequences. Thus the surjectivity of the map $(\Sigma_i f)_i$ follows from the surjectivity of $(f_{s(a)})_{t(a)=i}$ and f_i. Therefore $\Sigma_i f$ is surjective. $\square$

4.3. **Reflection functors and crystal operators $\mathcal{T}_i$.** We have seen in 2.6 that the operators $\mathcal{T}_i$ are analogues of Lusztig's braid group automorphisms T_i on crystal level. We will use the same notations $\mathcal{T}_i$ for their geometric counterpart on $\mathcal{B}$. An important result in [5] relates the reflection functors Σ_i^* (resp. Σ_i) to the crystal operators $\mathcal{T}_i$ (resp. $\mathcal{T}_i^{-1}$).

For any $b \in \mathscr{B}(\infty)$, denote its image of the isomorphism $\Psi : \mathscr{B}(\infty) \to \mathcal{B}$ by Z_b. Thus if $\mathrm{wt}(b) = -\nu$ for some $\nu \in Q_+$, $Z_b \in \mathrm{Irr}\,\Lambda(\nu)$.

Recall in 3.3 we have defined $\Lambda(\nu)_{i,c} = \{M \in \Lambda(\nu) | \dim \mathrm{top}_i M = c\}$. Similarly we define $\Lambda(\nu)_i^c = \{M \in \Lambda(\nu) | \dim \mathrm{soc}_i M = c\}$. The following results is the dual statement of [5] Proposition 18 (Note that our $\mathcal{T}_i$ (resp. ε_i, $\widetilde{e}_i$) was denoted by S_i^{-1} (resp. φ_i, $\widetilde{f}_i$) in [5]).

Proposition 4.4. *Let $b \in \mathscr{B}(\infty)$ with $\mathrm{wt}(b) = -\nu \in Q_-$. Set $d = \varepsilon_i^*(b)$, $U = Z_b \cap \Lambda(\nu)_i^c$, $b' = \mathcal{T}_i \widetilde{e}_i^{*\max} b$ and $\mathrm{wt}(b') = -\nu'$. Let V be an open dense $\mathrm{GL}(\nu')$-invariant subset of $Z_{b'}$. Then $\{M \in U | \mathcal{O}_{\Sigma_i^* M} \subseteq V\}$ contains a dense open subset of U.*

5. The modules $M_{\mathbf{i},k}$ via reflection functors

In this section we will show that the modules $M_{\mathbf{i},k}$ (recall 3.5) can be obtained from simple Λ-modules by applying reflection functors, which is crucial in the proof of our main results in this paper. In the proof we use the modules $N(w\varpi_i)$ defined in [5]. We show that they give an alternative construction of the modules $V_{\mathbf{i},k}$ (see 3.4). Using this result we can find the relationship between reflection functors and the modules $M_{\mathbf{i},k}$.

5.1. **Nakajima's quiver.** Let $\widehat{Q}$ be the quiver obtained from Q by adding a vertex i' and an arrow $d_i : i \to i'$ for each vertex $i \in I$. Then we have the associated preprojective algebra $\widehat{\Lambda} = \Lambda(\widehat{Q})$. So a $\widehat{\Lambda}$-module M consists of two I-graded vector spaces $\oplus_{i \in I} M_i$ and $\oplus_{i \in I} M_{i'}$ together with linear maps $M_a : M_{s(a)} \to M_{t(a)}$ (for each $a \in H$), $M_{d_i} : M_i \to M_{i'}$ and $M_{d_i^*} : M_{i'} \to M_i$ (for each $i \in I$) satisfying the following relations:

$$\sum_{a \in H, t(a)=i} \epsilon(a) M_a M_{a^*} = M_{d_i^*} M_{d_i}, \ M_{d_i} M_{d_i^*} = 0, \text{ for each } i \in I$$

It is convenience to write the dimension vector of a $\widehat{\Lambda}$-module M as a pair $(\mu, \lambda) \in Q_+ \times P_+$. That is, $\mu = \sum_{i \in I} \mu_i \alpha_i$ and $\lambda = \sum_{i \in I} \lambda_i \varpi_i$, where $\mu_i = \dim M_i$ and $\lambda_i = \dim M_{i'}$.

We say that a $\widehat{\Lambda}$-module M is *stable* if for each $i \in I$, $\operatorname{soc}_i M = 0$. That is, the socle of M is concentrated at primed vertices.

Remark 5.1. The above construction is slightly different from Nakajima's quiver varieties [19]. Namely in general the definition of Nakajima's quiver variety does not require the relation $M_{d_i} M_{d_i^*} = 0$. But the set of $\widehat{\Lambda}$-modules M with dimension vector (ν, λ) and $M_{d_i}^* = 0$ (for all $i \in I$) can be identified with the following variety

$$\Lambda(\nu, \lambda) = \Lambda(\nu) \times \bigoplus_{i \in I} \operatorname{Hom}_{\mathbb{C}}(M_i, M_{i'}),$$

And then the definition of stable $\widehat{\Lambda}$-modules coincides with the stability condition given in [19]. Denote the subset of stable points in $\Lambda(\nu, \lambda)$ by $\Lambda(\nu, \lambda)^{st}$. Nakajima defined a $\mathrm{GL}(\nu)$-action on $\Lambda(\nu, \lambda)$ and the stabilizer of any point in $\Lambda(\nu, \lambda)^{st}$ is trivial. So we have the variety $\mathcal{L}(\nu, \lambda) = \Lambda(\nu, \lambda)^{st} / \mathrm{GL}(\nu)$. This is a Lagrangian subvariety of Nakajima's quiver variety.

5.2. **The modules $\widehat{N}(w\lambda)$.** Let $\lambda \in P_+$ be a dominant weight. Define $\widehat{N}(\lambda)$ to be the $\widehat{\Lambda}$-module with dimension vector $(0, \lambda)$, which is unique up to isomorphism. it is clear that $\widehat{N}(\lambda)$ is stable.

For each $w \in W$, we define

$$\widehat{N}(w\lambda) = \Sigma_{i_r} \Sigma_{i_{r-1}} \cdots \Sigma_{i_1} \widehat{N}(\lambda).$$

where $\mathbf{i} = (i_r, \ldots, i_1)$ is a reduced expression of w. Note that by Lemma 4.2 (3), $\widehat{N}(w\lambda)$ is independent of the choice of $\mathbf{i}$ and is thus well-defined.

The following results are contained in [5] Theorem 8 and its proof there:

Lemma 5.2. *(1). $\widehat{N}(w\lambda)$ is the unique (up to isomorphism) stable $\widehat{\Lambda}$-module with dimension vector $(\lambda - w\lambda, \lambda)$.*

(2). For $i \in I$ such that $\ell(s_i w) > \ell(w)$, $\widehat{N}(w\lambda)$ has trivial i-top.

5.3. **The module $N(w\lambda)$.** For any module M we have a canonical morphism $M \to \Sigma_i M$, which can be seen from the following diagram

$$\begin{tikzcd}
\widetilde{M_i} \arrow[r, "M_{\mathrm{in}(i)}"] \arrow[d, "\mathrm{id}"] & M_i \arrow[r, "M_{\mathrm{out}(i)}"] \arrow[d, "\overline{M}_{\mathrm{out}(i)}"] & \widetilde{M_i} \arrow[d, "\mathrm{id}"] \\
\widetilde{M_i} \arrow[r, "\overline{M}_{\mathrm{out}(i)} M_{\mathrm{in}(i)}"] & \ker M_{\mathrm{in}(i)} \arrow[r, "\iota"] & \widetilde{M_i}
\end{tikzcd}$$

If M has trivial i-socle, which means that $M_{\mathrm{out}(i)}$ is injective, then from the diagram $\overline{M}_{\mathrm{out}(i)}$ is also injective. So the canonical morphism $M \to \Sigma_i M$ is a monomorphism.

Therefore we have the canonical embedding $\widehat{N}(w\lambda) \hookrightarrow \widehat{N}(s_i w\lambda)$ if $\ell(s_i w) > \ell(w)$. In particular, $\widehat{N}(\lambda)$ is a submodule of $\widehat{N}(w\lambda)$. We define $N(w\lambda)$ to be the quotient $\widehat{N}(w\lambda)/\widehat{N}(\lambda)$.

By the definition of reflection functors we know that applying Σ_i for some $i \in I$ to a $\widehat{\Lambda}$-module M will not change the vector spaces $M_{j'}$ for any j. Thus the underlying space of $\widehat{N}(\lambda)$ is exactly the sum of underlying spaces of $\widehat{N}(w\lambda)$ at primed vertices. Therefore $N(w\lambda)$ is in fact a Λ-module.

Remark 5.3. The existence of the inclusion $\widehat{N}(\lambda) \hookrightarrow \widehat{N}(w\lambda)$ forces the linear maps corresponding to arrows d_i^* in the module $\widehat{N}(w\lambda)$ to be zero for all $i \in I$. Therefore the module $\widehat{N}(w\lambda)$ lies in $\Lambda(\lambda - w\lambda, \lambda)^{st}$ (see Remark 5.1).

5.4. **An alternative construction of $V_{\mathbf{i},k}$.** From now on we fix a reduced expression $\mathbf{i} = (i_r, \dots, i_1)$. Recall that $V_{\mathbf{i},k} = \mathrm{soc}_{(i_k,\dots,i_1)} \widehat{I}_{i_k}$ (see 3.4). It was stated in [5] without proof that the modules $N(w\lambda)$ had been studied in [9]. In this subsection we give a precise statement (Proposition 5.5 below) and present a proof.

For a dominant weight $\lambda = \sum_{i \in I} \lambda_i \varpi_i$, we define an injective Λ-module $\widehat{I}^\lambda = \oplus_{i \in I} \widehat{I}_i^{\lambda_i}$.

Lemma 5.4. *For each $w \in W$, there is a unique (up to isomorphism) submodule of $\widehat{I}^\lambda$ with dimension vector $\lambda - w\lambda$.*

Proof. Let $Gr(\lambda - w\lambda, \widehat{I}^\lambda)$ be the projective variety consisting of all Λ-submodules of $\widehat{I}^\lambda$ with dimension vector $\lambda - w\lambda$. In [22] it was shown that the variety $Gr(\lambda - w\lambda, \widehat{I}^\lambda)$ is homeomorphic to the Lagrangian quiver variety $\mathcal{L}(\lambda - w\lambda, \lambda)$ (see Remark 5.1). Then the statement in the lemma is just a reformulation of Proposition 5.1 in [21], which asserts that the variety $\mathcal{L}(\lambda - w\lambda, \lambda)$ is a point. □

Now it is easy to prove the following result:

Proposition 5.5. *For any $1 \leq k \leq r$, $V_{\mathbf{i},k} \cong N(s_{i_1} s_{i_2} \cdots s_{i_k} \varpi_{i_k})$.*

Proof. In [5] Theorem 5, it was shown that $\mathrm{soc}\, N(w\varpi_i) \cong S_i$ and $\underline{\dim}\, N(w\varpi_i) = \varpi_i - w\varpi_i$ for any $w \neq 1 \in W$ and $i \in I$. In particular, the dimension vector of $N(s_{i_1} s_{i_2} \cdots s_{i_k} \varpi_{i_k})$ is $\varpi_{i_k} - s_{i_1} s_{i_2} \cdots s_{i_k} \varpi_{i_k}$. From [9] Corollary 9.2 we know that $V_{\mathbf{i},k}$ has the same dimension vector. Since both of them are submodules of $\widehat{I}_{i_k}$, they have to be isomorphic by the above lemma. □

5.5. **Reflection functors and the module $M_{\mathbf{i},k}$.** Now we are ready to prove the main result in this section. Recall that $M_{\mathbf{i},k} = V_{\mathbf{i},k} / V_{\mathbf{i},k^-}$.

Proposition 5.6. *For any $1 \leq k \leq r$, $M_{\mathbf{i},k} \cong \Sigma_{i_1} \Sigma_{i_2} \cdots \Sigma_{i_{k-1}} S_{i_k}$.*

Proof. First we have the following short exact sequence:

$$0 \to \widehat{N}(\varpi_{i_k}) \to \widehat{N}(s_{i_k} \varpi_{i_k}) \to S_{i_k} \to 0.$$

since $\widehat{N}(s_{i_k} \varpi_{i_k}) = \Sigma_{i_k} \widehat{N}(\varpi_{i_k})$ and $\widehat{N}(\varpi_{i_k}) = S_{i'_k}$.

By Lemma 4.2 (1), the functor Σ_i is left exact for any $i \in I$. Thus applying the functor $\Sigma_{i_{k-1}}$ we obtain the exact sequence

$$0 \to \Sigma_{i_{k-1}} \widehat{N}(\varpi_{i_k}) \to \Sigma_{i_{k-1}} \widehat{N}(s_{i_k} \varpi_{i_k}) \to \Sigma_{i_{k-1}} S_{i_k}.$$

But we know that $\mathrm{top}_{i_{k-1}} \widehat{N}(s_{i_k} \varpi_{i_k}) = 0$ by Lemma 5.2 (2). So Lemma 4.3 implies that the above sequence is in fact a short exact sequence.

We can repeat this procedure by applying $\Sigma_{i_{k-2}}, \dots, \Sigma_{i_1}$. Finally we have the following short exact sequence

$$0 \to \Sigma_{i_1} \cdots \Sigma_{i_{k-1}} \widehat{N}(\varpi_{i_k}) \to \Sigma_{i_1} \cdots \Sigma_{i_{k-1}} \widehat{N}(s_{i_k} \varpi_{i_k}) \to \Sigma_{i_1} \cdots \Sigma_{i_{k-1}} S_{i_k} \to 0. \tag{1}$$

Note that for any l such that $k^- < l < k$, we have $i_l \neq i_k$. This implies $\Sigma_{i_l} \widehat{N}(\varpi_{i_k}) = \widehat{N}(\varpi_{i_k})$ because the module $\widehat{N}(\varpi_{i_k})$ is concentrated at the vertex i'_k, which is not connected with any other vertex except i_k. So we have

$$\Sigma_{i_1} \Sigma_{i_2} \cdots \Sigma_{i_{k-1}} \widehat{N}(\varpi_{i_k}) = \Sigma_{i_1} \Sigma_{i_2} \cdots \Sigma_{i_{k^-}} \widehat{N}(\varpi_{i_k}).$$

Thus the above sequence (1) is the following

$$0 \to \widehat{N}(s_{i_1} s_{i_2} \cdots s_{i_{k^-}} \varpi_{i_k}) \to \widehat{N}(s_{i_1} s_{i_2} \cdots s_{i_k} \varpi_{i_k}) \to \Sigma_{i_1} \Sigma_{i_2} \cdots \Sigma_{i_{k-1}} S_{i_k} \to 0. \tag{2}$$

Now $\widehat{N}(s_{i_1} s_{i_2} \cdots s_{i_{k^-}} \varpi_{i_k})$ and $\widehat{N}(s_{i_1} s_{i_2} \cdots s_{i_k} \varpi_{i_k})$ both have the submodule $\hat{N}(\varpi_{i_k})$. And the map $\widehat{N}(s_{i_1} s_{i_2} \cdots s_{i_k} \varpi_{i_k}) \to \Sigma_{i_1} \Sigma_{i_2} \cdots \Sigma_{i_{k-1}} S_{i_k}$ clearly maps $\hat{N}(\varpi_{i_k})$ to zero. So sequence (2) yields

$$0 \to N(s_{i_1} s_{i_2} \cdots s_{i_{k^-}} \varpi_{i_k}) \to N(s_{i_1} s_{i_2} \cdots s_{i_k} \varpi_{i_k}) \to \Sigma_{i_1} \Sigma_{i_2} \cdots \Sigma_{i_{k-1}} S_{i_k} \to 0.$$

Applying Lemma 5.5 (note that $i_k = i_{k^-}$) we have

$$0 \to V_{\mathbf{i},k^-} \to V_{\mathbf{i},k} \to \Sigma_{i_1} \Sigma_{i_2} \cdots \Sigma_{i_{k-1}} S_{i_k} \to 0.$$

Hence $\Sigma_{i_1} \Sigma_{i_2} \cdots \Sigma_{i_{k-1}} S_{i_k} \cong V_{\mathbf{i},k} / V_{\mathbf{i},k^-} = M_{\mathbf{i},k}$. □

The following corollary will be needed in the next section:

Corollary 5.7. *For any $l < k-1$, the module $\Sigma_{i_{l+1}} \cdots \Sigma_{i_{k-1}} S_{i_k}$ has trivial i_l-top.*

Proof. In the proof of the above proposition we see that $\Sigma_{i_{l+1}} \cdots \Sigma_{k-1} S_{i_k}$ is a quotient of $\widehat{N}(s_{i_{l+1}} \cdots s_{i_{k-1}} \varpi_{i_k})$, which has trivial i_l-top by Lemma 5.2 (2). □

5.6. **The adaptable case.** Let Q be a quiver and $i \in I$ be a vertex, denote by $\sigma_i Q$ the quiver obtained from Q by reversing all the arrows connected with i, if i is a sink or a source of Q.

An element $w \in W$ is called *Q-adaptable* if there exists a reduced expression $\mathbf{i} = (i_r, \ldots, i_1)$ of $w \in W$ such that i_1 is a sink of Q and i_k is a sink of $\sigma_{i_{k-1}} \cdots \sigma_{i_i} Q$ for all $2 \leq k \leq r$. In this case we say that the reduced expression $\mathbf{i}$ is *Q-adapted.* An element $w \in W$ is called *adaptable* if there exists an orientation Q of the corresponding Dynkin graph such that w is Q-adaptable.

As pointed out in [9], if w is adaptable and $\mathbf{i}$ is Q^{op}-adapted, the module $M_{\mathbf{i}}$ is a *terminal $\mathbb{C}Q$-module* in the sense of [8]. This means that the modules $M_{\mathbf{i},k}$ ($1 \leq k \leq r$) are certain indecomposable preinjective $\mathbb{C}Q$-modules. In particular, in the case that $\mathfrak{g}$ is finite-dimensional, the longest element $w_0 \in W$ is always adaptable. If $\mathbf{i}$ is a Q^{op}-adapted reduced expression of w_0, the set $\{M_{\mathbf{i},k} | 1 \leq k \leq r\}$ forms a complete set of pairwise non-isomorphic indecomposable $\mathbb{C}Q$-modules. But in general there does not exist any quiver Q such that all the modules $M_{\mathbf{i},k}$ are $\mathbb{C}Q$-modules (see Example 5.8).

It is known that for any quiver Q, the indecomposable preprojective and preinjective modules can be obtained from simple modules via BGP-reflection functors (see [3] Theorem 1.3). Thus our Proposition 5.6 can be viewed as a generalization of this classical result to the case of any $w \in W$ and any reduced expression $\mathbf{i}$.

5.7. **Examples.** We conclude this section with some examples.

Example 5.8. Let Q be a quiver with underlying graph $1 \;\text{———}\; 2 \;\text{———}\; 3$. So Λ is the preprojecitve algebra of type A_3. Let $w = w_0 \in W$ the longest element and $\mathbf{i} = (2, 1, 2, 3, 2, 1)$.

First we calculate the modules $M_{\mathbf{i},k}$ by the original definition (see 3.4 and 3.5). For this we need to calculate $V_{\mathbf{i},k}$ first:

$$V_{\mathbf{i},1} = {\scriptstyle 1}\,, \qquad V_{\mathbf{i},2} = \begin{smallmatrix} 1 & \\ & 2 \end{smallmatrix}\,, \qquad V_{\mathbf{i},3} = \begin{smallmatrix} 1 & & \\ & 2 & \\ & & 3 \end{smallmatrix}$$

$$V_{\mathbf{i},4} = \begin{smallmatrix} 1 & & 3 \\ & 2 & \end{smallmatrix}\,, \qquad V_{\mathbf{i},5} = \begin{smallmatrix} & & 3 \\ & 2 & \\ 1 & & \end{smallmatrix}\,, \qquad V_{\mathbf{i},6} = \begin{smallmatrix} & 2 & \\ 1 & & 3 \\ & 2 & \end{smallmatrix}$$

Then we have:

$$M_{\mathbf{i},1} = V_{\mathbf{i},1} = \begin{smallmatrix}1\end{smallmatrix}, \qquad M_{\mathbf{i},2} = V_{\mathbf{i},2} = \begin{smallmatrix}1&\\&2\end{smallmatrix}, \qquad M_{\mathbf{i},3} = V_{\mathbf{i},3} = \begin{smallmatrix}1&&\\&2&\\&&3\end{smallmatrix}$$

$$M_{\mathbf{i},4} = V_{\mathbf{i},4}/V_{\mathbf{i},2} = \begin{smallmatrix}3\end{smallmatrix}, \qquad M_{\mathbf{i},5} = V_{\mathbf{i},5}/V_{\mathbf{i},1} = \begin{smallmatrix}&3\\2&\end{smallmatrix}, \qquad M_{\mathbf{i},6} = V_{\mathbf{i},6}/V_{\mathbf{i},4} = \begin{smallmatrix}2\end{smallmatrix}.$$

Next let's use Proposition 5.6 to calculate:

$$\begin{aligned}
M_{\mathbf{i},1} &= S_1 = \begin{smallmatrix}1\end{smallmatrix}\\
M_{\mathbf{i},2} &= \Sigma_1 S_2 = \begin{smallmatrix}1&\\&2\end{smallmatrix}\\
M_{\mathbf{i},3} &= \Sigma_1\Sigma_2 S_3 = \Sigma_1(\begin{smallmatrix}2&\\&3\end{smallmatrix}) = \begin{smallmatrix}1&&\\&2&\\&&3\end{smallmatrix}\\
M_{\mathbf{i},4} &= \Sigma_1\Sigma_2\Sigma_3 S_2 = \Sigma_1\Sigma_2(\begin{smallmatrix}&3\\2&\end{smallmatrix}) = \begin{smallmatrix}3\end{smallmatrix}\\
M_{\mathbf{i},5} &= \Sigma_1\Sigma_2\Sigma_3\Sigma_2 S_1 = \Sigma_1\Sigma_2(\begin{smallmatrix}&&3\\&2&\\1&&\end{smallmatrix}) = \Sigma_1(\begin{smallmatrix}&&3\\&2&\\1&&\end{smallmatrix}) = \begin{smallmatrix}&3\\2&\end{smallmatrix}\\
M_{\mathbf{i},6} &= \Sigma_1\Sigma_2\Sigma_3\Sigma_2\Sigma_1 S_2 = \Sigma_1\Sigma_2\Sigma_3 S_1 = \Sigma_1(\begin{smallmatrix}&2\\1&\end{smallmatrix}) = \begin{smallmatrix}2\end{smallmatrix}
\end{aligned}$$

We see that the results are the same.

Note that the reduced expression $\mathbf{i} = (2, 1, 2, 3, 2, 1)$ is not adapted to any Q (although $w = w_0$ is adaptable). It is easy to see that we cannot find any orientation Q of the graph $1 \,\rule[0.5ex]{2em}{0.4pt}\, 2 \,\rule[0.5ex]{2em}{0.4pt}\, 3$ such that all the modules $M_{\mathbf{i},k}$ are $\mathbb{C}Q$-modules.

Example 5.9. Again let Λ be the preprojecitve algebra of type A_3 and $w = w_0 \in W$ the longest element. This time we choose a Q-adapted reduced expression $\mathbf{i}' = (1, 2, 1, 3, 2, 1)$, where $Q = 1 \longleftarrow 2 \longleftarrow 3$. Similarly we can work out

$$\begin{aligned}
M_{\mathbf{i}',1} &= S_1 = \begin{smallmatrix}1\end{smallmatrix}\\
M_{\mathbf{i}',2} &= \Sigma_1 S_2 = \begin{smallmatrix}1&\\&2\end{smallmatrix}\\
M_{\mathbf{i}',3} &= \Sigma_1\Sigma_2 S_3 = \Sigma_1(\begin{smallmatrix}2&\\&3\end{smallmatrix}) = \begin{smallmatrix}1&&\\&2&\\&&3\end{smallmatrix}\\
M_{\mathbf{i}',4} &= \Sigma_1\Sigma_2\Sigma_3 S_1 = \Sigma_1\Sigma_2 S_1 = \Sigma_1(\begin{smallmatrix}&2\\1&\end{smallmatrix}) = \begin{smallmatrix}2\end{smallmatrix}\\
M_{\mathbf{i}',5} &= \Sigma_1\Sigma_2\Sigma_3\Sigma_1 S_2 = \Sigma_1\Sigma_2(\begin{smallmatrix}1&&3\\&2&\end{smallmatrix}) = \Sigma_1(\begin{smallmatrix}&2&\\1&&3\end{smallmatrix}) = \begin{smallmatrix}2&\\&3\end{smallmatrix}\\
M_{\mathbf{i}',6} &= \Sigma_1\Sigma_2\Sigma_3\Sigma_1\Sigma_2 S_1 = \Sigma_1\Sigma_2\Sigma_3 S_2 = \Sigma_1 S_3 = \begin{smallmatrix}3\end{smallmatrix}.
\end{aligned}$$

We can see that $\{M_{\mathbf{i}',k} | 1 \le k \le 6\}$ is the set of isomorphism classes of indecomposable representations of $Q^{\mathrm{op}} = 1 \longrightarrow 2 \longrightarrow 3$.

Example 5.10. Let Q be the quiver with underlying graph



So Λ is the preprojective algebra of type D_4. Let $w = s_1 s_4 s_3 s_2 s_4 s_3 s_2 s_1$ and $\mathbf{i} = (1, 4, 3, 2, 4, 3, 2, 1)$. It is easy to see that w is not adaptable. Let's calculate $M_{\mathbf{i},8}$.

First by definition:

$$V_{\mathbf{i},8} = \mathrm{soc}_{1,4,3,2,4,3,2,1}\, \widehat{I}_1 = \begin{smallmatrix}1&&\\&4&\\&2&3\\&4&\\1&&\end{smallmatrix}, \quad V_{\mathbf{i},8^-} = V_{\mathbf{i},1} = \begin{smallmatrix}1\end{smallmatrix}$$

Then

$$M_{\mathbf{i},8} = V_{\mathbf{i},8}/V_{\mathbf{i},1} = \begin{smallmatrix}1&&\\&4&\\&2&3\\&4&\end{smallmatrix}$$

Now by Proposition 5.6, $M_{\mathbf{i},8} = \Sigma_1\Sigma_2\Sigma_3\Sigma_4\Sigma_2\Sigma_3\Sigma_4 S_1$. We can verify it as following:

$$1 \xrightarrow{\Sigma_4} \begin{smallmatrix} & 4\\ 1 & \end{smallmatrix} \xrightarrow{\Sigma_3} \begin{smallmatrix} & & 3\\ & 4 & \\ 1 & & \end{smallmatrix} \xrightarrow{\Sigma_2} \begin{smallmatrix} & 2 & 3\\ & 4 & \\ 1 & & \end{smallmatrix} \xrightarrow{\Sigma_4} \begin{smallmatrix} & 4 & \\ & 2 & 3\\ & 4 & \\ 1 & & \end{smallmatrix} \xrightarrow{\Sigma_3} \begin{smallmatrix} & 4 & \\ & 2 & 3\\ & 4 & \\ 1 & & \end{smallmatrix} \xrightarrow{\Sigma_2} \begin{smallmatrix} & 4 & \\ & 2 & 3\\ & 4 & \\ 1 & & \end{smallmatrix} \xrightarrow{\Sigma_1} \begin{smallmatrix} 1 & & \\ & 4 & \\ & 2 & 3\\ & 4 & \end{smallmatrix}$$

6. Compatibility of two parametrizations

Throughout this section, let w be an element in the Weyl group and fix a reduced expression $\mathbf{i} = (i_r, \dots, i_1)$ of w.

6.1. **Main results.** Recall that we have injective maps $\psi_{\mathbf{i}} : \mathbb{N}^r \to \mathscr{B}(\infty)$ (see 2.7) and $\phi_{\mathbf{i}} : \mathbb{N}^r \to \mathcal{B}$ (see 3.5). Also we know the crystal isomorphism $\Psi : \mathscr{B}(\infty) \simeq \mathcal{B}$ (Theorem 3.2). So we have the following diagram:

$$\begin{tikzcd}
\mathbb{N}^r \arrow[r, "\psi_{\mathbf{i}}"] \arrow[dr, "\phi_{\mathbf{i}}"'] & \mathscr{B}(\infty) \arrow[d, "\Psi"] \\
 & \mathcal{B}
\end{tikzcd}$$

Our main result is that the above diagram is commutative. In other words, we have the following theorem:

Theorem 6.1. *$\Psi(b_{\mathbf{i},\mathbf{a}}) = Z_{\mathbf{i}}^{\mathbf{a}}$, for any $\mathbf{a} \in \mathbb{N}^r$.*

The theorem will be proved in 6.3. From the theorem it follows immediately that the subset $\mathcal{B}(w)$ is the geometric counterpart of the unipotent crystal $\mathscr{B}(w)$:

Corollary 6.2. *$\Psi(\mathscr{B}(w)) = \mathcal{B}(w)$.*

Remark 6.3. As mentioned in the introduction, in the case that $\mathfrak{g}$ is a finite-dimensional simple Lie algebra, $w = w_0$ the longest element in the Weyl group and $\mathbf{i}$ a Q-adapted reduced expression, the result in Theorem 6.1 has been proved in [5] Proposition 28.

In fact, in this case the set of modules $M_{\mathbf{i},k}$ $(1 \leq k \leq r)$ is exactly a complete set of pairwise non-isomorphic indecomposable representations of Q. For any $\mathbf{a} \in \mathbb{N}$, set $M_{\mathbf{i}}^{\mathbf{a}} = \oplus_{k=1}^{r} M_{\mathbf{i},k}^{a_k} \in \mathrm{mod}(\mathbb{C}Q)$. It is not difficult to see that $\Lambda_{\mathbf{i}}^{\mathbf{a}} = T^*\mathcal{O}_{M_{\mathbf{i}}^{\mathbf{a}}}$, the conormal bundle of the orbit of $M_{\mathbf{i}}^{\mathbf{a}}$. So the irreducible component $Z_{\mathbf{i}}^{\mathbf{a}}$ is the same as the closure of $T^*\mathcal{O}_{M_{\mathbf{i}}^{\mathbf{a}}}$.

The arguments in [5] also work for any Kac-Moody algebra $\mathfrak{g}$ and Q-adapted reduced expression $\mathbf{i}$ (which means that w is Q- adaptable). However, even in the case of finite type, there exists non-adaptable w in the Weyl group (see Example 5.10). Thus our theorem is new in call cases.

Remark 6.4. In general, for any $b \in \mathscr{B}(\infty)$, the image $\Psi(b) \in \mathcal{B}$ is not easy to describe. We can only use the fact that Ψ commutes with $\widetilde{e}_i$, $\widetilde{f}_i$ and keeps the unique highest weight element. More precisely, denote by b_∞ (resp. Z_0) the unique highest weight element in $\mathscr{B}(\infty)$ (resp. $\mathcal{B}$). One need to find a path in the crystal graph from b_∞ to b, say, $b = \widetilde{f}_{j_1}\widetilde{f}_{j_2}\cdots\widetilde{f}_{j_s}b_\infty$. Then we have $\Psi(b) = \widetilde{f}_{j_1}\widetilde{f}_{j_2}\cdots\widetilde{f}_{j_s}Z_0$.

We see that Theorem 6.1 gives a direct description of the image $\Psi(b)$ for any $b \in \mathscr{B}(w)$ once we know the $\mathbf{i}$-Lusztig data of b for some $\mathbf{i}$.

6.2. **Calculation of Lusztig data.** In this subsection we make some preparations to prove the main theorem.

The following proposition shows how to deduce the $\mathbf{i}$-Lusztig data $(\psi_{\mathbf{i}}^{-1}(b))$ of any $b \in \mathscr{B}(w)$ by applying operators $\widetilde{e}_i^*$ and $\mathcal{T}_i$ (recall 2.6). It was essentially proved in [16] and [20].

Proposition 6.5. *Let* $b \in \mathscr{B}(w)$ *and assume that* $\psi_{\mathbf{i}}^{-1}(b) = \mathbf{a} = (a_r, \dots, a_1)$. *Then*

(1). $a_1 = \varepsilon_{i_1}^*(b)$. $\psi_{\mathbf{i}}^{-1}(\widetilde{e}_i^{*\max} b) = (a_r, \dots, a_2, 0)$.

(2). Let $w' = s_{i_r} \cdots s_{i_2}$ *and denote by* $\mathbf{i}' = (i_r, \dots, i_2)$, $\mathbf{a}' = (a_r, \dots, a_2)$. *Suppose that* $a_1 = 0$, *then* $\mathcal{T}_{i_1}(b) \in \mathscr{B}(w')$ *and we have* $\psi_{\mathbf{i}'}^{-1}(\mathcal{T}_{i_1}(b)) = \mathbf{a}'$.

Proof. We just recall the proof of (2) here. Remember $G : \mathscr{B}(\infty) \to \mathbf{B}$ is the globalization map from the crystal basis to the canonical basis.

Since $a_1 = 0$, we have

$$G(b) \equiv F_{\mathbf{i}}(\mathbf{a}) = T_{i_1}^{-1} \cdots T_{i_{r-1}}^{-1}(f_{i_r}^{(a_r)}) \cdots T_{i_1}^{-1}(f_{i_2}^{(a_2)}), \quad \mod q\mathscr{L}(\infty).$$

Thus

$$T_{i_1}(G(b)) \equiv T_{i_2}^{-1} \cdots T_{i_{r-1}}^{-1}(f_{i_r}^{(a_r)}) \cdots T_{i_2}^{-1}(f_{i_3}^{(a_3)}) f_{i_2}^{(a_2)} \equiv F_{\mathbf{i}'}(\mathbf{a}').$$

By Proposition 2.1, $T_{i_1}(G(b)) \equiv \mathcal{T}_{i_1}(b)$. This ends the proof. □

Combine (1) and (2) in the above proposition we have $\mathcal{T}_{i_1} \widetilde{e}_{i_1}^{*\max} b_{\mathbf{i},\mathbf{a}} = b_{\mathbf{i}',\mathbf{a}'}$. We can repeat the procedure and finally we will reach the unique highest weight element b_∞. This gives the following corollary.

Corollary 6.6. $\widetilde{e}_{i_r}^{*\max} \mathcal{T}_{i_{r-1}} \widetilde{e}_{i_{r-1}}^{*\max} \cdots \mathcal{T}_{i_1} \widetilde{e}_{i_1}^{*\max} b_{\mathbf{i},\mathbf{a}} = b_\infty$.

6.3. **Proof of theorem 6.1.** Now we are ready to prove the main theorem.

As in Proposition 6.5 we denote by $\mathbf{i}' = (i_r, \dots, i_2)$ and $\mathbf{a}' = (a_r, \dots, a_2)$.

Lemma 6.7. *For any* $X \in \Lambda_{\mathbf{i}}^{\mathbf{a}}$, $\Sigma_{i_1}^* X \in \Lambda_{\mathbf{i}'}^{\mathbf{a}'}$.

Proof. By definition X has a filtration

$$0 = X_0 \subseteq X_1 \subseteq \cdots \subseteq X_r = X \tag{3}$$

such that $X_k / X_{k-1} \cong M_{\mathbf{i},k}^{a_k}$ for any $1 \leq k \leq r$. Denote by $Y = X/X_1$. We have the following exact sequence

$$0 \to X_1 \to X \to Y \to 0.$$

Applying $\Sigma_{i_1}^*$ to the above sequence, we have $\Sigma_{i_1}^* X \cong \Sigma_{i_1}^* Y$ because $X_1 \cong M_{\mathbf{i},1}^{a_1} = S_{i_1}^{a_1}$, $\Sigma_{i_1}^* S_{i_1} = 0$ and the functor $\Sigma_{i_1}^*$ is right exact (Lemma 4.2).

The filtration (3) induces the following filtration of Y:

$$0 = Y_1 \subseteq Y_2 \subseteq \cdots \subseteq Y_r = Y,$$

where $Y_k \cong X_k / X_1$ for any $2 \leq k \leq r$. In particular, $Y_k / Y_{k-1} \cong X_k / X_{k-1} \cong M_{\mathbf{i},k}^{a_k}$.

By Proposition 5.6, $M_{\mathbf{i},k} = \Sigma_{i_1} \cdots \Sigma_{i_{k-1}} S_{i_k}$ for any $1 \leq k \leq r$. So for $2 \leq k \leq r$, $M_{\mathbf{i},k}$ has trivial i_1-socle. Then by Lemma 4.3, we know that $\Sigma_{i_1}^* Y$ has a filtration

$$0 = Y_1' \subseteq Y_2' \subseteq \cdots \subseteq Y_r' = Y$$

such that $Y_k' / Y_{k-1}' \cong (\Sigma_{i_1}^* M_{\mathbf{i},k})^{a_k}$.

Note that $\Sigma_{i_2} \cdots \Sigma_{i_{k-1}} S_{i_k}$ has trivial i_1-top by Corollary 5.7. By Lemma 4.2 (2), we deduce that

$$\Sigma_{i_1}^* M_{\mathbf{i},k} = \Sigma_{i_1}^* \Sigma_{i_1} \Sigma_{i_2} \cdots \Sigma_{i_{k-1}} S_{i_k} \cong \Sigma_{i_2} \cdots \Sigma_{i_{k-1}} S_{i_k}.$$

This proves $\Sigma_{i_1}^* Y \in \Lambda_{\mathbf{i}'}^{\mathbf{a}'}$. □

Lemma 6.8. $\mathcal{T}_{i_1}\widetilde{e}^{*\,\max}_{i_1}(Z^{\mathbf{a}}_{\mathbf{i}}) = Z^{\mathbf{a}'}_{\mathbf{i}'}$.

Proof. For any $X \in \Lambda^{\mathbf{a}}_{\mathbf{i}}$, since $M_{\mathbf{i},k}$ has trivial i_1-socle for all $2 \le k \le r$, we have

$$\operatorname{soc}_{i_1} X \cong \operatorname{soc}_{i_1} M_{\mathbf{i},i_1} = S^{a_1}_{i_1}.$$

We know that the irreducible component $Z^{\mathbf{a}}_{\mathbf{i}}$ contains a dense open subset belonging to $\Lambda^{\mathbf{a}}_{\mathbf{i}}$, thus $\varepsilon^*_{i_1}(Z^{\mathbf{a}}_{\mathbf{i}}) = a_1$.

Now by Proposition 4.4, there exists a dense open subset U of $Z^{\mathbf{a}}_{\mathbf{i}}$ such that $U \subseteq \Lambda^{\mathbf{a}}_{\mathbf{i}}$ and for any $X \in U$, $\Sigma^*_{i_1}X$ lies in a dense open subset of $\mathcal{T}_{i_1}\widetilde{e}^{*\,\max}_{i_1}(Z^{\mathbf{a}}_{\mathbf{i}})$.

So Lemma 6.7 implies $\mathcal{T}_{i_1}\widetilde{e}^{*\,\max}_{i_1}(Z^{\mathbf{a}}_{\mathbf{i}}) = Z^{\mathbf{a}'}_{\mathbf{i}'}$. □

We can apply Lemma 6.8 successively to $Z^{\mathbf{a}}_{\mathbf{i}}$ until we reach the unique irreducible component of the variety $\Lambda(0)$ (in fact it is a point), which is the highest weight element Z_0 in $\mathcal{B}$. Namely we have

$$(**) \qquad \widetilde{e}^{*\,\max}_{i_r}\mathcal{T}_{i_{r-1}}\widetilde{e}^{*\,\max}_{i_{r-1}}\cdots\mathcal{T}_{i_1}\widetilde{e}^{*\,\max}_{i_1}Z^{\mathbf{a}}_{\mathbf{i}} = Z_0.$$

Now just note that the crystal isomorphism $\Psi : \mathscr{B}(\infty) \simeq \mathcal{B}$ maps b_∞ to Z_0 and commutes with $\widetilde{e}^*_i$ and $\mathcal{T}_i$. Comparing $(**)$ with Corollary 6.6, we complete the proof of Theorem 6.1.

7. Interpretations of Lusztig's transition map $R^{\mathbf{i}'}_{\mathbf{i}}$

7.1. **The transition map.** For two different reduced expressions $\mathbf{i}, \mathbf{i}'$ of a Weyl group element w, the parametrizations $\phi_{\mathbf{i}}, \phi_{\mathbf{i}'} : \mathbb{N}^r \to \mathscr{B}(w)$ are in general different. So we have the transition map $R^{\mathbf{i}'}_{\mathbf{i}} : \mathbb{N}^r \to \mathbb{N}^r$ defined by

$$R^{\mathbf{i}'}_{\mathbf{i}} = \phi^{-1}_{\mathbf{i}'} \circ \phi_{\mathbf{i}}.$$

Lusztig [16] has described the transition map $R^{\mathbf{i}'}_{\mathbf{i}}$ as a composition of some simple piecewise-linear transformations. Let's recall his description here.

For a reduced expression $\mathbf{i}$, a 2-*move* on $\mathbf{i}$ is to replace two consecutive entries (i,j) by (j,i) when $c_{ij} = 0$. A 3-*move* is to replace three consecutive entries (i,j,i) by (j,i,j) when $c_{ij} = -1$. It is known that any two reduced expressions of an element $w \in W$ can be obtained from each other by a sequence of 2-moves and 3-moves. Thus we only need to describe the transition map $R^{\mathbf{i}'}_{\mathbf{i}}$ when $\mathbf{i}$ and $\mathbf{i}'$ differ by a 2-move or a 3-move. Lusztig's descriptions are the following:

Proposition 7.1. *For* $\mathbf{a} \in \mathbb{N}^r$, *let* $\mathbf{a}' = R^{\mathbf{i}'}_{\mathbf{i}}(\mathbf{a})$.

1. *If* $\mathbf{i}$ *and* $\mathbf{i}'$ *differ by a 2-move, i.e. for some* k, $i'_k = i_{k+1}$, $i'_{k+1} = i_k$ *and* $i'_l = i_l$ *for all* $l \neq k, k+1$, *Then* $a'_k = a_{k+1}$, $a'_{k+1} = a_k$ *and* $a'_l = a_l$ *for all* $l \neq k, k+1$.
2. *If* $\mathbf{i}$ *and* $\mathbf{i}'$ *differ by a 3-move, i.e. for some* k, $i'_{k+1} = i'_{k-1} = i_k$, $i'_k = i_{k+1} = i_{k-1}$ *and* $i'_l = i_l$ *for all* $l \neq k-1, k, k+1$. *Then*

$$\begin{aligned} a'_{k+1} &= a_k + a_{k-1} - \min(a_{k+1}, a_{k-1}),\\ a'_k &= \min(a_{k+1}, a_{k-1}),\\ a'_{k-1} &= a_k + a_{k+1} - \min(a_{k+1}, a_{k-1}) \end{aligned}$$

   and $a'_l = a_l$ *for all* $l \neq k-1, k, k+1$.

The following corollary follows immediately from Theorem 6.1:

Corollary 7.2. $Z^{\mathbf{a}}_{\mathbf{i}} = Z^{\mathbf{a}'}_{\mathbf{i}'}$ *if and only if* $R^{\mathbf{i}'}_{\mathbf{i}}(\mathbf{a}) = \mathbf{a}'$.

Recall that $Z_{\mathbf{i}}^{\mathbf{a}}$ is the Zariski closure of the constructible set $\Lambda_{\mathbf{i}}^{\mathbf{a}}$. So there exists an $X \in \Lambda_{\mathbf{i}}^{\mathbf{a}} \cap \Lambda_{\mathbf{i}'}^{\mathbf{a}'}$ in a dense open subset of $Z_{\mathbf{i}}^{\mathbf{a}}$ such that the parameter $\mathbf{a}$ (resp. $\mathbf{a}'$) encodes the multiplicities of the modules $M_{\mathbf{i},s}$ (resp. $M_{\mathbf{i}',s}$) in certain filtrations of X. On the other hand, for any $X \in \Lambda_{\mathbf{i}}^{\mathbf{a}}$, we know that there exists a unique $\mathbf{b} \in \mathbb{N}$ such that $X \in \Lambda_{\mathbf{i}'}^{\mathbf{b}}$ (see 3.5). We will see that in general $\mathbf{b} \neq R_{\mathbf{i}}^{\mathbf{i}'}(\mathbf{a})$.

In the following we focus on the cases that $\mathbf{i}$ and $\mathbf{i}'$ differ by a 2-move or a 3-move. First we find out the difference between the two sets of modules $M_{\mathbf{i},s}$ and $M_{\mathbf{i}',s}$ $(1 \leq s \leq r)$. Then we try to explain how the parameter $\mathbf{a}$ changes to $\mathbf{b}$ for $X \in \Lambda_{\mathbf{i}}^{\mathbf{a}} \cap \Lambda_{\mathbf{i}'}^{\mathbf{b}}$. In particular, it gives interpretations of the transition map $R_{\mathbf{i}}^{\mathbf{i}'}$ when X is in a dense open subset of $Z_{\mathbf{i}}^{\mathbf{a}}$.

7.2. **The case of 2-moves.** As before let $\mathbf{i} = (i_r, \ldots, i_1)$ be a reduced expression of $w \in W$. First we deduce an easy lemma:

Lemma 7.3. *For $1 \leq s \leq r$ and $s^- < k \leq s$, we have $\mathrm{Ext}^1_\Lambda(M_{\mathbf{i},s}, M_{\mathbf{i},k}) = 0$.*

Proof. Apply $\mathrm{Hom}_\Lambda(-, M_{\mathbf{i},k})$ to the exact sequence:

$$0 \to V_{\mathbf{i},s^-} \to V_{\mathbf{i},s} \to M_{\mathbf{i},s} \to 0,$$

we have the following exact sequence

$$\mathrm{Hom}_\Lambda(V_{\mathbf{i},s^-}, M_{\mathbf{i},k}) \to \mathrm{Ext}^1_\Lambda(M_{\mathbf{i},s}, M_{\mathbf{i},k}) \to \mathrm{Ext}^1_\Lambda(V_{\mathbf{i},s}, M_{\mathbf{i},k}).$$

Since $M_{\mathbf{i},k} = V_{\mathbf{i},k}/V_{\mathbf{i},k^-}$ lies in the subcategory $\mathcal{C}_{(i_s,\cdots,i_1)}$ for $k \leq s$ and $V_{\mathbf{i},s}$ is $\mathcal{C}_{(i_s,\cdots,i_1)}$-projective-injective, we know that $\mathrm{Ext}^1_\Lambda(V_{\mathbf{i},s}, M_{\mathbf{i},k}) = 0$. From [9] Lemma 9.8 (1) we know that $\mathrm{Hom}_\Lambda(V_{\mathbf{i},s^-}, M_{\mathbf{i},k}) = 0$ for any $k > s^-$. Thus $\mathrm{Ext}^1_\Lambda(M_{\mathbf{i},s}, M_{\mathbf{i},k}) = 0$ for any $s^- < k \leq s$. □

Now we assume that $\mathbf{i}$ and $\mathbf{i}'$ are two reduced expressions of $w \in W$ differing by a 2-move. More precisely, there exists $1 \leq k < r$ such that $i'_k = i_{k+1}$, $i'_{k+1} = i_k$, $i'_l = i_l$ for all $l \neq k, k+1$ and $c_{i_k i_{k+1}} = 0$ in the Cartan matrix. For simplicity we will write $V_k = V_{\mathbf{i},k}$, $M_k = M_{\mathbf{i},k}$, $V'_k = V_{\mathbf{i}',k}$ and $M'_k = V_{\mathbf{i}',k}$ in the sequel.

We can see from the definition that $V'_{k+1} = V_k$, $V'_k = V_{k+1}$. Therefore $M'_{k+1} = M_k$ and $M'_k = M_{k+1}$. So the set of modules $\{M_s | 1 \leq s \leq r\}$ is invariant when we change parametrizations. Only two of them change their index. Now we can prove the following result:

Proposition 7.4. *With the notations above, we have $\Lambda_{\mathbf{i}}^{\mathbf{a}} = \Lambda_{\mathbf{i}'}^{\mathbf{a}'}$, where $\mathbf{a}' = R_{\mathbf{i}}^{\mathbf{i}'}(\mathbf{a}')$.*

Proof. Let X be a module in $\Lambda_{\mathbf{i}}^{\mathbf{a}}$. So X has a filtration

$$0 = X_0 \subseteq X_1 \subseteq \cdots \subseteq X_r = X$$

such that $X_s/X_{s-1} \simeq M_s^{a_s}$ for all $1 \leq s \leq r$.

By Lemma 7.3, $\mathrm{Ext}^1_\Lambda(M_k, M_{k+1}) = 0$. This means that we can exchange M_k and M_{k+1} in the above filtration of X, which yields a new filtration

$$0 = X'_0 \subseteq X'_1 \subseteq \cdots \subseteq X'_r = X$$

where $X'_s = X_s$ for $s \neq k$, $X'_k/X'_{k-1} \simeq M_{k+1}^{a_{k+1}}$ and $X'_{k+1}/X'_k \simeq M_k^{a_k}$.

Now by Proposition 7.1, we have $a'_k = a_{k+1}$, $a'_{k+1} = a_k$. Thus the above filtration of X satisfies $X'_s/X'_{s-1} \cong M_s'^{a'_s}$ for any $1 \leq s \leq r$, which implies $X \in \Lambda_{\mathbf{i}'}^{\mathbf{a}'}$. □

Thus we see that in this case the change of parameters is quite simple, namely it satisfies Lusztig's transition map for all the modules $X \in \Lambda_{\mathbf{i}}^{\mathbf{a}}$, not only those in a dense subset of $Z_{\mathbf{i}}^{\mathbf{a}}$.

7.3. **The case of 3-moves.** In this subsection we deal with the case that $\mathbf{i}$ and $\mathbf{i}'$ differ by a 3-move. So we assume that for some k, $i'_{k+1} = i'_{k-1} = i_k$, $i'_k = i_{k+1} = i_{k-1}$ and $i'_l = i_l$ for all $l \neq k-1, k, k+1$. And $c_{i_k i_{k+1}} = -1$ in the Cartan matrix. As in the last subsection we write V_k (resp. V'_k) for $V_{\mathbf{i},k}$ (resp. $V_{\mathbf{i}',k}$) and M_k (resp. M'_k) for $M_{\mathbf{i},k}$ (resp. $M_{\mathbf{i}',k}$).

Lemma 7.5. $\dim \operatorname{Ext}^1_\Lambda(M_{k+1}, M_{k-1}) = \dim \operatorname{Ext}^1_\Lambda(M_{k-1}, M_{k+1}) = 1.$

Proof. The first equality is just Lemma 3.1. We prove the second one.

By Proposition 5.6 we have,

$$M_{k-1} = \Sigma_{i_1} \cdots \Sigma_{i_{k-2}}(S_{i_{k-1}}),\ M_{k+1} = \Sigma_{i_1} \cdots \Sigma_{i_k}(S_{i_{k+1}}).$$

Then using Corollary 5.7 and Lemma 4.3 we can deduce that

$$\operatorname{Ext}^1_\Lambda(M_{k+1}, M_{k-1}) = \operatorname{Ext}^1_\Lambda(\Sigma_{i_{k-1}}\Sigma_{i_k}(S_{i_{k+1}}), S_{i_{k-1}}).$$

Note that $i_{k-1} = i_{k+1}$ and $c_{i_k i_{k+1}} = -1$. It is easy to see that

$$\Sigma_{i_{k-1}}\Sigma_{i_k}(S_{i_{k+1}}) = S_{i_k}.$$

So we have

$$\dim \operatorname{Ext}^1_\Lambda(M_{k+1}, M_{k-1}) = \dim \operatorname{Ext}^1_\Lambda(S_{i_k}, S_{i_{k-1}}) = 1.$$

□

According to the lemma, there is a unique (up to isomorphism) non-trivial extension of M_{k-1} by M_{k+1}. And so is the extension of M_{k+1} by M_{k-1}.

The following results describe the difference between the modules M_s and M'_s for all $1 \leq s \leq r$.

Proposition 7.6. (1) *$M_l = M'_l$ for $l \neq k-1, k, k+1$.*

(2) *We have $M'_{k-1} = M_{k+1}$, $M'_{k+1} = M_{k-1}$ and M'_k is the unique non-trivial extension of M_{k+1} by M_{k-1} (so M_k is the unique non-trivial extension of M_{k-1} by M_{k+1}).*

Proof. By [4] Lemma III 4.2, we know that the maximal rigid module $V_{\mathbf{i}'} = \oplus_{k=1}^r V'_k$ can be obtained from $V_{\mathbf{i}} = \oplus_{k=1}^r V_k$ by a single step of mutation. More precisely, we have $V'_{k+1} = V_k$, $V'_k = V_{k+1}$ and $V'_{k-1} = V^*_{k-1}$, all other V_s $(s \neq k-1, k, k+1)$ remains unchanged. The first statement of the proposition follows immediately. We refer to [4] or [9] for all missing definitions and notations about maximal rigid modules and their mutations.

It is not difficult to see that the two exchange sequences are as follows:

$$0 \to V_{k-1} \to V_{k^-} \oplus V_{k+1} \to V^*_{k-1} \to 0,$$

$$0 \to V^*_{k-1} \to V_k \oplus V_{(k-1)^-} \to V_{k-1} \to 0.$$

For each module $X \in \Lambda^{\mathbf{a}}_{\mathbf{i}}$, denote by $M_{\mathbf{i}}(X) = \oplus_{k=1}^r M_k^{a_k}$. The first sequence above is $\operatorname{Hom}_\Lambda(V_{\mathbf{i}}, -)$-exact, hence is $M_{\mathbf{i}}$-split ([9], Corollary 10.7), which means that

$$M_{\mathbf{i}}(V_{k-1}) \oplus M_{\mathbf{i}}(V^*_{k-1}) = M_{\mathbf{i}}(V_{k^-}) \oplus M_{\mathbf{i}}(V_{k+1}).$$

Note that $M_{\mathbf{i}}(V_{k+1}) = M_{\mathbf{i}}(V_{k-1}) \oplus M_{k+1}$. So splitting out $M_{\mathbf{i}}(V_{k-1})$ from both sides we have

$$M_{\mathbf{i}}(V^*_{k-1}) = M_{\mathbf{i}}(V_{k^-}) \oplus M_{k+1}. \tag{4}$$

By the definition of M'_{k-1} we have the exact sequence

$$0 \to V'_{(k-1)^-} \to V'_{k-1} \to M'_{k-1} \to 0,$$

which is also $\mathrm{Hom}_\Lambda(V_{\mathbf{i}}, -)$-exact since $V'_{(k-1)^-} = V_{k^-}$. Thus we have

$$M_{\mathbf{i}}(V^*_{k-1}) = M_{\mathbf{i}}(V'_{k-1}) = M_{\mathbf{i}}(V_{k^-}) \oplus M_{\mathbf{i}}(M'_{k-1}). \tag{5}$$

Compare (5) with (4), we have $M_{\mathbf{i}}(M'_{k-1}) = M_{k+1}$. Therefore $M'_{k-1} = M_{k+1}$. Similarly $M'_{k+1} = M_{k-1}$.

By the definition of M'_k we have the exact sequence

$$0 \to V'_{k^-} \to V'_k \to M'_k \to 0,$$

Since $V'_{k^-} = V_{(k-1)^-}$, the sequence is also $M_{\mathbf{i}}$-split. Note that $V'_k = V_{k+1}$. So we have

$$M_{\mathbf{i}}(V_{k+1}) = M_{\mathbf{i}}(V_{(k-1)^-}) \oplus M_{\mathbf{i}}(M'_k).$$

But we also have

$$M_{\mathbf{i}}(V_{k+1}) = M_{\mathbf{i}}(V_{(k-1)^-}) \oplus M_{k-1} \oplus M_{k+1}.$$

So we can deduce that $M_{\mathbf{i}}(M'_k) = M_{k-1} \oplus M_{k+1}$.

We know that M'_k is indecomposable. And we already proved $M'_{k-1} = M_{k+1}$ and $M'_{k+1} = M_{k-1}$. By the fact that $\mathcal{C}_{\mathbf{i},\mathbf{a}}$ ($\mathbf{a} \in \mathbb{N}$) are pairwise disjoint (3.5), the only possibility is that M'_k is the unique extension of M_{k+1} by M_{k-1}. Similarly M_k is the unique extension of M_{k-1} by M_{k+1}. ☐

Example 7.7. Let Λ be the preprojective algebra of type A_3. $\mathbf{i} = (2, 1, 2, 3, 2, 1)$ and $\mathbf{i}' = (1, 2, 1, 3, 2, 1)$. So $\mathbf{i}$ and $\mathbf{i}'$ differ by a 3-move and $k = 5$.

From Example 5.8 and Example 5.9 we see that

$$M_1 = M'_1 = {\scriptstyle 1}, \quad M_2 = M'_2 = \begin{smallmatrix} 1 & \\ & 2 \end{smallmatrix}, \quad M_3 = M'_3 = \begin{smallmatrix} 1 & & \\ & 2 & \\ & & 3 \end{smallmatrix}$$

and

$$\begin{array}{lll} M_4 = {\scriptstyle 3}, & M_5 = \begin{smallmatrix} & 3 \\ 2 & \end{smallmatrix}, & M_6 = {\scriptstyle 2} \\ M'_4 = {\scriptstyle 2}, & M'_5 = \begin{smallmatrix} 2 & \\ & 3 \end{smallmatrix}, & M'_6 = {\scriptstyle 3} \end{array}$$

whose relationship is exactly as described in the above proposition.

Thus we see that the set of modules $\{M_s | 1 \leq s \leq r\}$ does not change too much when $\mathbf{i}$ is replaced by $\mathbf{i}'$. In fact only one of them, namely M_k changes. And the two modules M_{k+1}, M_{k-1} change their indices. Moreover the M_k, which is the extension of M_{k-1} by M_{k+1}, is just replaced by the extension module in the other direction. In particular, there are only two "building blocks" in the three modules, namely M_{k+1} and M_{k-1}. Therefore we can expect that different parametrizations correspond to different arrangement of these two "building blocks". In the following we try to explain it in details.

Let $X \in \Lambda^{\mathbf{a}}_{\mathbf{i}}$ (not necessarily be in a dense open subset of $Z^{\mathbf{a}}_{\mathbf{i}}$), X has a filtration

$$0 \subseteq X_0 \subseteq X_1 \subseteq \cdots \subseteq X_r = X \tag{6}$$

such that $X_s/X_{s-1} \cong M_s^{a_s}$ for any $1 \leq s \leq r$.

Now we focus on the part X_{k+1}/X_{k-2}. By Lemma 7.3, we have

$$\mathrm{Ext}^1_\Lambda(M_{k-1}, M_k) = \mathrm{Ext}^1_\Lambda(M_k, M_{k+1}) = 0.$$

So the only possible indecomposable direct summands of X_{k+1}/X_{k-2} are M_{k-1}, M_{k+1} and their extensions. We may visualize X_{k+1}/X_{k-2} as follows:

$$\begin{array}{cccccccc} M_{k-1} & & M_{k+1} & & & & & \\ | & \cdots & | & \cdots & M_{k-1} & \cdots & M_{k+1} & \cdots \\ M_{k+1} & & M_{k-1} & & & & & \\ (M_k) & & (M'_k) & & & & & \end{array} \tag{7}$$

Suppose that $X_{k+1}/X_{k-2} = M_k^{n_1} \oplus M_k'^{n_2} \oplus M_{k-1}^{n_3} \oplus M_{k+1}^{n_4}$, $n_j \in \mathbb{N}$ for $1 \leq j \leq 4$. Define $\Lambda_{\mathbf{i}}^{\mathbf{a}}(k; n_1, n_2, n_3, n_4)$ to be the subset of $\Lambda_{\mathbf{i}}^{\mathbf{a}}$ consisting of those X with a filtration (6), where $X_{k+1}/X_{k-2} \simeq M_k^{n_1} \oplus M_k'^{n_2} \oplus M_{k-1}^{n_3} \oplus M_{k+1}^{n_4}$.

It is clear that:

$$a_{k-1} = n_2 + n_3, \quad a_k = n_1, \quad a_{k+1} = n_2 + n_4.$$

We have already mentioned that there is a unique $\mathbf{b} \in \mathbb{N}^r$ such that $X \in \Lambda_{\mathbf{i}'}^{\mathbf{b}}$. The above picture (7) shows that $X \in \Lambda_{\mathbf{i}'}^{\mathbf{b}}(k; n_2, n_1, n_4, n_3)$, so we have

$$b_{k-1} = n_1 + n_4, \quad b_k = n_2, \quad b_{k+1} = n_1 + n_3.$$

Unfortunately we are not able to express (b_{k-1}, b_k, b_{k+1}) with (a_{k-1}, a_k, a_{k+1}) directly. But if $\min\{n_3, n_4\} = 0$ we can do it and it is as follows:

$$\begin{aligned} b_{k+1} &= a_k + a_{k-1} - \min(a_{k+1}, a_{k-1}), \\ b_k &= \min(a_{k+1}, a_{k-1}), \\ b_{k-1} &= a_k + a_{k+1} - \min(a_{k+1}, a_{k-1}) \end{aligned}$$

which is exactly Lusztig's transition map (Proposition 7.1). Thus we have proved the following

Proposition 7.8. *For any $X \in \Lambda_{\mathbf{i}}^{\mathbf{a}}(k; n_1, n_2, n_3, n_4)$, if $X \in \Lambda_{\mathbf{i}'}^{\mathbf{b}}$, then $\mathbf{b} = R_{\mathbf{i}}^{\mathbf{i}'}(\mathbf{a})$ if and only if $\min\{n_3, n_4\} = 0$.*

We give a simple example showing the case that $\mathbf{b} \neq R_{\mathbf{i}}^{\mathbf{i}'}(\mathbf{a})$.

Example 7.9. Let Λ be the preprojective algebra of type A_2. Let $\mathbf{i} = (1, 2, 1)$, $\mathbf{i}' = (2, 1, 2)$ be two reduced expressions of the longest Weyl group element. We have $M_1 = {}_1$, $M_2 = {}^1{}_2$, $M_3 = {}_2$, and $M'_1 = {}_2$, $M'_2 = {}^2{}_1$, $M'_3 = {}_1$.

Now consider the module $X = {}^1{}_2 \oplus {}_1 \oplus {}_2$. It is obvious that X is in $\Lambda_{\mathbf{i}}^{(1,1,1)}$ and $\Lambda_{\mathbf{i}'}^{(2,0,2)}$. However, by Proposition 7.1 we have $R_{\mathbf{i}}^{\mathbf{i}'}(1, 1, 1) = (1, 1, 1)$.

Thus we know that in general $\Lambda_{\mathbf{i}}^{\mathbf{a}} \neq \Lambda_{\mathbf{i}}^{\mathbf{a}'}$, where $\mathbf{a}' = R_{\mathbf{i}}^{\mathbf{i}'}(\mathbf{a})$. However, we also know by Corollary 7.2 that $Z_{\mathbf{i}}^{\mathbf{a}} = Z_{\mathbf{i}'}^{\mathbf{a}'}$. It means that there exists a dense open subset U of $Z_{\mathbf{i}}^{\mathbf{a}}$ belonging to $\Lambda_{\mathbf{i}}^{\mathbf{a}} \cap \Lambda_{\mathbf{i}}^{\mathbf{a}'}$. Thus for any $X \in U$, if $X \in \Lambda_{\mathbf{i}}^{\mathbf{a}}(k; n_1, n_2, n_3, n_4)$, then the condition $\min\{n_3, n_4\} = 0$ must be satisfied.

Acknowledgments. Part of this work was done when the author was visiting the Hausdorff Research Institute for Mathematics (HIM) in Bonn as a participant of the trimester program "On the interaction of representation theory with geometry and combinatorics". The author would like to thank the organizers for their hospitality. He is also grateful to Prof. Jan Schröer for interesting discussions and helpful comments.

References

1. C. Amiot, O. Iyama, I. Reiten and G. Todorov, Preprojective algebras and c-sortable words, Preprint arXiv:1002.4131.
2. A. Berenstein, S. Fomin and A. Zelevinsky, Parametrizations of canonical bases and totally positive matrices, Adv. Math. 122(1996), 49-149.
3. I. N. Berstein, I. M. Gelfand and V. A. Ponomarev, Coxeter functors and Gabriel's theorem, *Uspehi. Mat. Nauk* 28 (1973), 19-33.
4. A. Buan, O. Iyama, I. Reiten and J. Scott. Cluster structures for 2-Calabi-Yau categories and unipotent groups, *Compositio Math.* 145 (2009), 1035-1079.
5. P. Baumann and J. Kamnitzer, Preprojective algebras and MV polytopes, Preprint arXiv:1009.2469.
6. A. Berenstein and A. Zelevinsky, Tensor product multiplicities, canonical bases and totally positive varieties, *Invent. Math.* 143(2001), 77-128.
7. W. Crawley-Boevey, On the exeptional fibres of Kleinian singularities, *Amer. J. Math.* 122 (2000), 1027-1037.
8. C. Geiß, B. Leclerc and J. Schröer, Cluster algerba structures and semicanonical bases for unipotent groups, Preprint arXiv:0703039.
9. C. Geiß, B. Leclerc and J. Schröer, Kac-Moody groups and cluster algebras, *Adv. Math.* 228 (2011), 329-433.
10. M. Kashiwara, On crystal bases of the q-anlogue of universal enveloping algebras, *Duke Math. J.* 63 (1991), 456-516.
11. M. Kashiwara, The crystal base and Littelmann's refined Demazure character formula, *Duke Math. J.* 71 (1993), 839-858.
12. M. Kashiwara and Y. Saito, Geometric construction of crystal bases, *Duke Math. J.* 89 (1997), 9-36.
13. Y. Kimura, Quantum unipotent subgroups and dual canonical bases, Preprint arXiv:1010.4212.
14. G. Lusztig, Canonical bases arising from quantized enveloping algebras, *J. Amer. Math. Soc.* 3 (1990), 447-498.
15. G. Lusztig, Quivers, pervere sheaves and quantized enveloping algebras, *J. Amer. Math. Soc.* 4 (1991), 365-421.
16. G. Lusztig, Introduction to quantum groups, Birkhäuser, Boston, 1993.
17. G. Lusztig, Braid group actions and canonical bases, *Adv. Math.* 122 (1996), 237-261.
18. G. Lusztig, Semicanonical bases arising from enveloping algebras, *Adv. Math.* 151 (2000), 129-139.
19. H. Nakajima, Instantons on ALE spaces, quiver varities and Kac-Moody algebras, *Duke Math. J.* 76 (1994), 365-416.
20. Y. Saito, PBW-basis of quantized universal enveloping algebras, *Publ. RIMS, Kyoto Univ.* 30 (1994), 209-232.
21. A. Savage, Quiver varieties and Demazure modules, *Math. Ann.* 355 (2006), 31-46.
22. A. Savage, Quiver grassmannians, quiver vareties and the preprojective algebra, *Pacific J. Math.* 251 (2011), 393-430.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, POSTFACH 10 01 31, D-33501 BIELEFELD, GERMANY
E-mail address: `yjiang@math.uni-bielefeld.de`